

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 August 2001 (16.08.2001)

PCT

(10) International Publication Number
WO 01/58945 A1

(51) International Patent Classification⁷: **C07K 14/47**,
14/705, 14/72, 16/18, 16/22, 16/26, G01N 33/53, 33/567,
C12N 5/10, 15/12, 15/63, 15/64

(74) Agent: **WALLEN, John, W., III**; Johnson & Johnson,
One Johnson & Johnson Plaza, New Brunswick, NJ 08933
(US).

(21) International Application Number: PCT/US01/03456

(22) International Filing Date: 1 February 2001 (01.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/500,123 8 February 2000 (08.02.2000) US

(71) Applicant: **ORTHO-MCNEIL PHARMACEUTICAL, INC.** [US/US]; U.S. Route No. 202, Raritan, NJ 08869 (US).

(72) Inventors: **DUBIN, Adrienne, Elizabeth**; 4303 Bromfield Avenue, San Diego, CA 92122 (US). **HUVAR, Arne**; 5560 Shasta Lane #67, La Mesa, CA 91942 (US). **GLASS, Charles, A.**; 10454 Woodchuck Point, San Diego, CA 92131 (US). **ERLANDER, Mark, G.**; 442 Hillcrest, Encinitas, CA 92024 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DNA ENCODING HUMAN VANILLOID RECEPTOR VR3

(57) Abstract: DNA encoding human VR1 receptor has been cloned and characterized. The recombinant protein is capable of forming biologically active protein. The cDNA's have been expressed in recombinant host cells that produce active recombinant protein. The recombinant protein is also purified from the recombinant host cells. In addition, the recombinant host cells are utilized to establish a method for identifying modulators of the receptor activity, and receptor modulators are identified.



WO 01/58945 A1

DNA ENCODING HUMAN VANILLOID RECEPTOR VR3

BACKGROUND OF THE INVENTION

Noxious chemical, thermal and mechanical stimuli excite peripheral nerve endings of small diameter sensory neurons (nociceptors) in sensory ganglia (eg., dorsal root, nodose and trigeminal ganglia) and initiate signals that are perceived as pain. These neurons are crucial for the detection of harmful or potentially harmful stimuli (heat) and tissue damage (H^+ (local tissue acidosis), and/or stretch) which arise from changes in the extracellular space during inflammatory or ischaemic conditions (Wall and Melzack, 1994). The vanilloid capsaicin (8-methyl-N-vanillyl-6-nonenamide), the main pungent ingredient in “hot” capsicum peppers, is a very selective activator of thinly or unmyelinated nociceptive afferents (Szolcsanyi, 1993; Szolcsanyi, 1996). Electrophysiological studies have shown that vanilloids excite small sensory neurons by activating a plasma membrane channel that is non-selectively permeable to cations (Bevan and Szolcsanyi, 1990; Oh et al., 1996; Wood et al., 1988). The ultra potent tricyclic diterpene resiniferatoxin from *Euphorbia* plants (RTX; (Szolcsanyi et al., 1991)) binds with nanomolar affinity at the capsaicin binding site and has revealed a very localized distribution of capsaicin receptors to rat somatic and visceral primary sensory neurons (Szallasi, 1995).

The vanilloid receptor VR1 (Caterina et al., 1997) is thought to be a heat-sensing receptor whose threshold is decreased in the presence of protons or capsaicin (Tominaga et al., 1998). Capsaicin and protons interact at specific membrane recognition sites (vanilloid receptors) expressed almost exclusively by primary sensory neurons involved in nociception and neurogenic inflammation (Bevan and Szolcsanyi, 1990). The vanilloid (“capsaicin”) receptor VR1 is activated by capsaicin and RTX, and activation of VR1 is blocked by the antagonists capsazepine (CPZ; (Bevan et al., 1992)) and ruthenium red (RR; (Caterina et al., 1997; Wood et al., 1988)).

Hydropathicity analysis of the amino acid sequence of VR1 reveals 6 potential membrane spanning regions (S1-S6) and a putative pore-loop region between S5 and S6. A large intracellular domain contains 3 ankyrin repeat domains. (Caterina et al., 1997). This channel has significant structural similarities with the putative “store-operated” TRP calcium channel family. VR1 is a ligand-gated non-selective cation channel that shows pronounced outward rectification (Caterina et al., 1997). Importantly, VR1 is highly permeable to Ca^{2+} , an ion known to be very important in regulating cell function ((Blackstone and Sheng, 1999; Gupta and Pushkala, 1999; van Haasteren et al., 1999)).

Searching genomic databases has revealed VRL-1, a subunit structurally related to VR1. Rat and human VRL-1 (AF129113 and AF129112, respectively) are ~49% identical and 66% similar to rat VR1 (AF029310)) (Caterina et al., 1999). Human VRL protein (AF103906) cloned by Wood and colleagues (unpublished) is 99% identical to VRL-1 (AF129112). Recently, a patent application by Partiseti and Renard was published that described hVRCC (human vanilloid receptor like cation channel) which is nearly identical to AF129112 (the only difference is the deletion of Q418). We will refer to these sequences as VR2. Overall, the predicted structure of VR1 and VR2 is characteristic of a family of ion channels defined by the transient receptor potential (TRP) channels originally cloned from *Drosophila melanogaster*, a Ca^{2+} -permeable channel that plays a role in phototransduction (Lu and Wong, 1987; Minke and Selinger, 1996). This receptor appears to also be involved in the sensation of pain-producing heat (Caterina et al., 1999). Expression of VR2 in oocytes and HEK cells usually conferred a sensitivity of the cells to noxious temperatures ($>53^\circ\text{C}$), that was not sensitive to CPZ but was nearly completely blocked at $10\ \mu\text{M}$ ruthenium red. Activation of VR2 induces a non-selective cation current with high permeability to Ca^{2+} . Interestingly, the threshold for heat sensitivity decreased with repeated application of noxious stimuli, but not subthreshold temperatures (Caterina et al., 1999).

The rat SIC (stretch-inhibitable channel; Genbank AB015231), encoded by 529 amino acids, is thought to form an ion channel inhibited by stretch (Suzuki et al., 1999). The first 379 amino acids homologous to rat VR1. SIC lacks the large N-terminal cytoplasmic domain of the VR family but contains a sequence homologous to the A exon prior to the putative TM1. The last 163 amino acids, beginning in the middle of putative TM6 of rat SIC are similar to the corresponding amino acid sequence of the human VR3 A+B- of the present invention.

The present invention describes the cloning and function of a novel vanilloid receptor family member, VR3. This gene appears to be alternatively spliced to create at least 3 isoforms.

SUMMARY OF THE INVENTION

DNA molecules encoding 3 isoforms of the human vanilloid receptor 3 (hVR3) have been cloned and characterized. The biological and structural properties of these proteins are disclosed, as is the amino acid and nucleotide sequence. The recombinant protein is useful to identify modulators of the receptor VR3. Modulators identified in the assay disclosed herein are useful as therapeutic agents, which are candidates for the treatment of inflammatory conditions and for use as analgesics for intractable pain associated with postherpetic neuralgia, diabetic neuropathy, postmastectomy pain, complex regional pain syndromes, arthritis (e.g., rheumatoid and osteoarthritis), as well as ulcers, neurodegenerative diseases, asthma, chronic obstructive pulmonary disease, irritable bowel syndrome, and psoriasis. Uses include the treatment of central nervous system diseases, diseases of the intestinal tract, abnormal proliferation and cancer especially in the digestive system, prostate and female gonads, ulcer, liver disease, kidney disease, control of viscera innervated by the dorsal root ganglia, or to diagnose or treat any disorder related to abnormal expression of these hVR3 polypeptides, among others. In another aspect, the

invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with hVR3 imbalance. The recombinant DNA molecules, and portions thereof, are useful for isolating homologues of the DNA molecules, identifying and isolating genomic equivalents of the DNA molecules, and identifying, detecting or isolating mutant forms of the DNA molecules.

BRIEF DESCRIPTION OF THE DRAWING

FIGURE 1 - SEQ.ID.NO.:5. Human VR3A+B- nucleotide sequence of the coding region (2616 bp).

FIGURE 2 - SEQ.ID.NO.:6. The nucleotide sequence of human VR3A+B- is shown including 337 bp 5' untranslated region (UT) and 547 bp 3'UT (3500 bp).

FIGURE 3 - SEQ.ID.NO.:7. Coding sequence for human VR3A+B- (871 amino acids)

FIGURE 4 - SEQ.ID.NO.:8. Human VR3A-B- nucleotide sequence of the coding region (2436 bp).

FIGURE 5 - SEQ.ID.NO.:9. Coding sequence for human VR3A-B- (811 amino acids)

FIGURE 6 - SEQ.ID.NO.:10. Human VR3A+B+ nucleotide sequence of the coding region (2229 bp).

FIGURE 7 - SEQ.ID.NO.:11. The nucleotide sequence of human VR3A+B+ is shown including 836 bp 5' UT and 994 bp 3'UT (4059 bp).

FIGURE 8 - SEQ.ID.NO.:12. Coding sequence for human VR3A+B+ (742 amino acids)

FIGURE 9 – Functional expression of VR3 isoforms in *Xenopus* oocytes is shown: viability of oocyte maintained in ND-96 with 2 Ca^{2+} was significantly diminished 4-6 days after injection with 3.25 ng VR3 A+B- and VR3 A+B+ (5 ng) but not VR3 A-B- (1.5 ng) cRNA. Data for VR3 A+B- and water-injected oocytes were obtained from 5 different experiments. Dead oocytes were determined visually. Data were analyzed using Chi square analysis.

FIGURE 10 - Function in oocytes: VR3 isoforms are activated by heat. A. Shown is the mean peak current elicited by a heat ramp from 25 deg C to 46 deg C and maintained at 46 degC for at least 15 sec. The current is the increase over initial current at +80 mV. Voltage ramps were applied from -120 to +80 mV over 400 msec every 2 sec. Oocytes were injected with 3.25, 1.5 and 5 ng VR3 A+B-, A-B-, and A+B+ cRNA, respectively, and recorded up to 6 days later. Solid bar: water injected controls (n=8); clear bar: VR3 A+B- isoform (n= 11); hatched bar: VR3 A-B- isoform (n= 8); stippled bar: VR3 A+B+ isoform (n=5). All isoforms show a significant increase over water controls (p= 0.001, 0.03 and 0.007, respectively; Student's *t*-test). The heat induced response is about 2-fold larger in the A+B- isoform

compared to the other 2 isoforms but the differences are not significant ($p=0.051$ and $p=0.16$, for A+B- compared to A-B- and A+B+, respectively). Data were obtained from 2 sets of injected oocytes. Data shown is the mean and standard error of the mean. B. Voltage-ramp induced currents were recorded during application of increasing heat to oocytes injected with water (top), VR3 A+B- (second from top), VR3 A-B- (3rd from top), and VR3A+B+ (bottom). Oocytes were constantly perfused with Ca²⁺ ND-96 and the solution was heated by an inline heater device (TC-324B in conjunction with the SH-27A in line heater; Warner Instrument Corp.). Ramp induced currents (in uA, as indicated on the y-axis) obtained at temperatures from 37 deg C to 46 deg C are displayed; only current traces at the higher temperatures are labeled with the corresponding temperature.

FIGURE 11 - Function in oocytes: VR3A+B- confers sensitivity to 10 uM ruthenium red to the perfusion activated current ($I_{\text{perfusion}}$) observed in VR1-expressing oocytes induced by cell perfusion by extracellular saline solutions. Activation of $I_{\text{perfusion}}$ by increased perfusion was blocked by ruthenium red only in VR1-expressing oocytes injected with VR3 A+B- cRNA, and not in oocytes expressing only VR1. (a). An oocyte injected with VR1 cRNA [4 ng] was challenged with voltage ramps between -120 and +80 mV over 400 msec from a holding potential of -70 mV. The ramp-induced currents were increased after onset of perfusion of the oocyte at a rate of 10 ml/min. Preincubation with 10 uM ruthenium red for 0.5 min blocked the current. The block was partially reversible (not shown). (b). An oocyte

injected with VR1 cRNA [4 ng] together with VR3 A+B- [1.3 ng] was challenged with voltage ramps between -120 and $+80$ mV over 400 msec from a holding potential of -70 mV. The ramp-induced currents were increased after onset of perfusion of the oocyte at a rate of 10 ml/min. Preincubation with 10 μ M ruthenium red for 0.5 min had little effect on $I_{\text{perfusion}}$. $I_{\text{perfusion}}$ had similar magnitudes in both sets of oocytes. V_{rev} for the RR inhibited current was about -13 mV (arrow).

FIGURE 12 – Function in oocytes: VR3A+B- together with VR1. The magnitude and decay kinetics of the response to 1 μ M capsaicin was diminished when VR1 cRNA was co-expressed with VR3 A+B- cRNA at equal ratios [2.9 ng each].

FIGURE 13- DNA array distribution analysis indicates that hVR3 mRNAs are expressed in a variety of tissues, and there is some overlap of expression with VR1 at a whole tissue level. The DNA sequence used on the DNA array is not present in A+B+, only A+B- and A-B- isoforms. Note that the cDNA species was cloned from pituitary and prostate glands.

DETAILED DESCRIPTION

The present invention describes 3 isoforms of a human vanilloid receptor, termed VR3: VR3A+B-, VR3A-B-, and VR3A+B+. The nucleotide sequences of human VR3 receptor cDNAs revealed single large open reading frame of about 2616 (Figure 1), 2436 (Figure 4) and 2229 (Figure 6) base pairs encoding 871 (Figure 3), 811 (Figure 5), and 742 (Figure 8) amino acids for human VR3 A+B-, A-B- and

A+B+, respectively. The cDNA for VR3 A+B- has 5' and 3'-untranslated extensions of about 337 and about 547 nucleotides, as shown in Figure 2, wherein the 5'UTR is 1-337, the coding region is 338 – 2953, and the 3'UTR is 2954 - 3500. The cDNA for VR3 A+B+ has 5' and 3'-untranslated extensions of about 836 and about 994 nucleotides as shown in Figure 7, wherein the 5'UTR is 1-836, the coding region is 837 – 3065, and the 3'UTR is 3066 - 4059. The first in-frame methionine was designated as the initiation codon for an open reading frame that predicts human VR3 receptor proteins with an estimated molecular mass (M_r) of about 98,242 Da, 91,294 Da and 83,310 Da for the isoforms A+B-, A-B- and A+B+, respectively. The A+B- isoform encodes a protein of 871 amino acids. The VR3 A-B- contains a deletion of 60 amino acids from amino acid 382 to 441 of VR3A+B-. VR3 A+B+ is identical to A+B- until amino acid 736 after which there are 6 divergent amino acids and a stop codon. The VR3 A+B+ isoform extends 20 amino acids after the putative TM6.

The predicted human VR3 receptor proteins were aligned with nucleotide and protein databases and are related to the vanilloid receptor family (VR1 and VR2). There are several conserved motifs found in this family of receptor including a large putative N-terminal hydrophilic segment (about 467 amino acids), three putative ankyrin repeat domains in the N-terminus region, 6 predicted transmembrane regions and a pore region. VR3 A+B- is 43 % identical to human VR1, 39% identical to both human VRL-1 (AF129112) and human VRL (AF103906). Thus the VR3 receptor described herein is clearly a novel gene of the vanilloid receptor family.

Human VR3A+B- and VR3A-B- forms are similar to the rat stretch-inhibitable channel SIC [Genbank accession AB015231] from amino acid 694 to the end. Rat SIC, encoded by 529 amino acids, is thought to form an ion channel inhibited by stretch. It lacks the large N-terminal cytoplasmic domain of the VR family but contains a sequence homologous to the A exon prior to the putative TM1.

The complete genomic sequence of the VR3 coding regions described herein appears to be found in a 380512 base pair sequence submission to Genbank (homo

sapiens clone RPC11-7G5 (AC007834), direct submission by Worley, K.C.). This Genbank entry lists many fragments of DNA sequence and a proposed contiguous sequence, but lacks any analysis of the nucleic acid sequence and fails to characterize the features of the VR3 nucleic acid sequences, or describe the presence of the VR3 gene.

Isolation of human VR3 receptor nucleic acid

The present invention relates to DNA encoding human VR3 receptor which were isolated from human VR3 receptor producing cells. Human VR3 receptor, as used herein, refers to protein which can specifically function as a human vanilloid receptor.

The complete amino acid sequence of human VR3 receptor was not previously known, nor was the complete nucleotide sequence encoding human VR3 receptor known. It is predicted that a wide variety of cells and cell types will contain the described human VR3 receptor.

Other cells and cell lines may also be suitable for use to isolate human VR3 receptor cDNA. Selection of suitable cells may be done by screening for human VR3 receptor activity in cell extracts or in whole cell assays, described herein. Cells that possess human VR3 receptor activity in any one of these assays may be suitable for the isolation of human VR3 receptor DNA or mRNA.

Any of a variety of procedures known in the art may be used to molecularly clone human VR3 receptor DNA. These methods include, but are not limited to, direct functional expression of the human VR3 receptor genes following the construction of a human VR3 receptor-containing cDNA library in an appropriate expression vector system. Another method is to screen human VR3 receptor-containing cDNA library constructed in a bacteriophage or plasmid shuttle vector with a labelled oligonucleotide probe designed from the amino acid sequence of the human VR3 receptor subunits. An additional method consists of screening a human VR3 receptor-containing cDNA library constructed in a bacteriophage or plasmid

shuttle vector with a partial cDNA encoding the human VR3 receptor protein. This partial cDNA is obtained by the specific PCR amplification of human VR3 receptor DNA fragments through the design of degenerate oligonucleotide primers from the amino acid sequence of the purified human VR3 receptor protein.

Another method is to isolate RNA from human VR3 receptor-producing cells and translate the RNA into protein via an *in vitro* or an *in vivo* translation system. The translation of the RNA into a peptide a protein will result in the production of at least a portion of the human VR3 receptor protein which can be identified by, for example, immunological reactivity with an anti-human VR3 receptor antibody or by biological activity of human VR3 receptor protein. In this method, pools of RNA isolated from human VR3 receptor-producing cells can be analyzed for the presence of an RNA that encodes at least a portion of the human VR3 receptor protein. Further fractionation of the RNA pool can be done to purify the human VR3 receptor RNA from non-human VR3 receptor RNA. The peptide or protein produced by this method may be analyzed to provide amino acid sequences which in turn are used to provide primers for production of human VR3 receptor cDNA, or the RNA used for translation can be analyzed to provide nucleotide sequences encoding human VR3 receptor and produce probes for this production of human VR3 receptor cDNA. This method is known in the art and can be found in, for example, Maniatis, T., Fritsch, E.F., Sambrook, J. in Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 1989.

It is readily apparent to those skilled in the art that other types of libraries, as well as libraries constructed from other cells or cell types, may be useful for isolating human VR3 receptor-encoding DNA. Other types of libraries include, but are not limited to, cDNA libraries derived from other cells, from organisms other than human VR3 receptor, and genomic DNA libraries that include YAC (yeast artificial chromosome) and cosmid libraries.

It is readily apparent to those skilled in the art that suitable cDNA libraries may be prepared from cells or cell lines which have human VR3 receptor activity. The selection of cells or cell lines for use in preparing a cDNA library to isolate human VR3 receptor cDNA may be done by first measuring cell associated human VR3 receptor activity using the measurement of human VR3 receptor-associated biological activity or a ligand binding assay.

Preparation of cDNA libraries can be performed by standard techniques well known in the art. Well known cDNA library construction techniques can be found for example, in Maniatis, T., Fritsch, E.F., Sambrook, J., *Molecular Cloning: A Laboratory Manual*, Second Edition (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989).

It is also readily apparent to those skilled in the art that DNA encoding human VR3 receptor may also be isolated from a suitable genomic DNA library. Construction of genomic DNA libraries can be performed by standard techniques well known in the art. Well known genomic DNA library construction techniques can be found in Maniatis, T., Fritsch, E.F., Sambrook, J. in *Molecular Cloning: A Laboratory Manual*, Second Edition (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989).

In order to clone the human VR3 receptor gene by the above methods, the amino acid sequence of human VR3 receptor may be necessary. To accomplish this, human VR3 receptor protein may be purified and partial amino acid sequence determined by automated sequenators. It is not necessary to determine the entire amino acid sequence, but the linear sequence of two regions of 6 to 8 amino acids from the protein is determined for the production of primers for PCR amplification of a partial human VR3 receptor DNA fragment.

Once suitable amino acid sequences have been identified, the DNA sequences capable of encoding them are synthesized. Because the genetic code is degenerate, more than one codon may be used to encode a particular amino acid, and therefore,

the amino acid sequence can be encoded by any of a set of similar DNA oligonucleotides. Only one member of the set will be identical to the human VR3 receptor sequence but will be capable of hybridizing to human VR3 receptor DNA even in the presence of DNA oligonucleotides with mismatches. The mismatched DNA oligonucleotides may still sufficiently hybridize to the human VR3 receptor DNA to permit identification and isolation of human VR3 receptor encoding DNA. DNA isolated by these methods can be used to screen DNA libraries from a variety of cell types, from invertebrate and vertebrate sources, and to isolate homologous genes.

Purified biologically active human VR3 receptor may have several different physical forms. human VR3 receptor may exist as a full-length nascent or unprocessed polypeptide, or as partially processed polypeptides or combinations of processed polypeptides. The full-length nascent human VR3 receptor polypeptide may be posttranslationally modified by specific proteolytic cleavage events that results in the formation of fragments of the full length nascent polypeptide. A fragment, or physical association of fragments may have the full biological activity associated with human VR3 receptor however, the degree of human VR3 receptor activity may vary between individual human VR3 receptor fragments and physically associated human VR3 receptor polypeptide fragments.

Because the genetic code is degenerate, more than one codon may be used to encode a particular amino acid, and therefore, the amino acid sequence can be encoded by any of a set of similar DNA oligonucleotides. Only one member of the set will be identical to the human VR3 receptor sequence but will be capable of hybridizing to human VR3 receptor DNA even in the presence of DNA oligonucleotides with mismatches under appropriate conditions. Under alternate conditions, the mismatched DNA oligonucleotides may still hybridize to the human VR3 receptor DNA to permit identification and isolation of human VR3 receptor encoding DNA.

DNA encoding human VR3 receptor from a particular organism may be used to isolate and purify homologues of human VR3 receptor from other organisms. To accomplish this, the first human VR3 receptor DNA may be mixed with a sample containing DNA encoding homologues of human VR3 receptor under appropriate hybridization conditions. The hybridized DNA complex may be isolated and the DNA encoding the homologous DNA may be purified therefrom.

It is known that there is a substantial amount of redundancy in the various codons that code for specific amino acids. Therefore, this invention is also directed to those DNA sequences that contain alternative codons that code for the eventual translation of the identical amino acid. For purposes of this specification, a sequence bearing one or more replaced codons will be defined as a degenerate variation. Also included within the scope of this invention are mutations either in the DNA sequence or the translated protein that do not substantially alter the ultimate physical properties of the expressed protein. For example, substitution of valine for leucine, arginine for lysine, or asparagine for glutamine may not cause a change in functionality of the polypeptide. Such substitutions are well known and are described, for instance in Molecular Biology of the Gene, 4th Ed. Benjamin Cummings Pub. Co. by Watson *et al.*

It is known that DNA sequences coding for a peptide may be altered so as to code for a peptide having properties that are different than those of the naturally occurring peptide. Methods of altering the DNA sequences include, but are not limited to site directed mutagenesis, chimeric substitution, and gene fusions. Site-directed mutagenesis is used to change one or more DNA residues that may result in a silent mutation, a conservative mutation, or a nonconservative mutation. Chimeric genes are prepared by swapping domains of similar or different genes to replace similar domains in the human VR3 receptor gene. Similarly, fusion genes may be prepared that add domains to the human VR3 receptor gene, such as an affinity tag to facilitate identification and isolation of the gene. Fusion genes may be prepared to

replace regions of the human VR3 receptor gene, for example to create a soluble version of the protein by removing a transmembrane domain or adding a targeting sequence to redirect the normal transport of the protein, or adding new post-translational modification sequences to the human VR3 receptor gene. Examples of altered properties include but are not limited to changes in the affinity of an enzyme for a substrate or a receptor for a ligand.

As used herein, a "functional derivative" of human VR3 receptor is a compound that possesses a biological activity (either functional or structural) that is substantially similar to the biological activity of human VR3 receptor. The term "functional derivatives" is intended to include the "fragments," "variants," "degenerate variants," "analogs" and "homologues" or to "chemical derivatives" of human VR3 receptor. The term "fragment" is meant to refer to any polypeptide subset of human VR3 receptor. The term "variant" is meant to refer to a molecule substantially similar in structure and function to either the entire human VR3 receptor molecule or to a fragment thereof. A molecule is "substantially similar" to human VR3 receptor if both molecules have substantially similar structures or if both molecules possess similar biological activity. Therefore, if the two molecules possess substantially similar activity, they are considered to be variants even if the structure of one of the molecules is not found in the other or even if the two amino acid sequences are not identical. The term "analog" refers to a molecule substantially similar in function to either the entire human VR3 receptor molecule or to a fragment thereof.

Recombinant expression of human VR3 receptor

The cloned human VR3 receptor DNA obtained through the methods described herein may be recombinantly expressed by molecular cloning into an expression vector containing a suitable promoter and other appropriate transcription regulatory elements, and transferred into prokaryotic or eukaryotic host cells to

produce recombinant human VR3 receptor protein. Techniques for such manipulations are fully described in Maniatis, T, *et al.*, *supra*, and are well known in the art.

Expression vectors are defined herein as DNA sequences that are required for the transcription of cloned copies of genes and the translation of their mRNAs in an appropriate host. Such vectors can be used to express eukaryotic genes in a variety of hosts such as bacteria including *E. coli*, blue-green algae, plant cells, insect cells, fungal cells including yeast cells, and animal cells.

Specifically designed vectors allow the shuttling of DNA between hosts such as bacteria-yeast or bacteria-animal cells or bacteria-fungal cells or bacteria-invertebrate cells. An appropriately constructed expression vector should contain: an origin of replication for autonomous replication in host cells, selectable markers, a limited number of useful restriction enzyme sites, a potential for high copy number, and active promoters. A promoter is defined as a DNA sequence that directs RNA polymerase to bind to DNA and initiate RNA synthesis. A strong promoter is one that causes mRNAs to be initiated at high frequency. Expression vectors may include, but are not limited to, cloning vectors, modified cloning vectors, specifically designed plasmids or viruses.

A variety of mammalian expression vectors may be used to express recombinant Human VR3 receptor in mammalian cells. Commercially available mammalian expression vectors which may be suitable for recombinant Human VR3 receptor expression, include but are not limited to, pMAMneo (Clontech), pcDNA3 (InVitrogen), pMC1neo (Stratagene), pXT1 (Stratagene), pSG5 (Stratagene), EBO-pSV2-neo (ATCC 37593) pBPV-1(8-2) (ATCC 37110), pdBPV-MMTneo(342-12) (ATCC 37224), pRSVgpt (ATCC 37199), pRSVneo (ATCC 37198), pSV2-dhfr (ATCC 37146), pUCTag (ATCC 37460), and IZD35 (ATCC 37565).

A variety of bacterial expression vectors may be used to express recombinant human VR3 receptor in bacterial cells. Commercially available bacterial expression

vectors which may be suitable for recombinant human VR3 receptor expression include, but are not limited to pET vectors (Novagen) and pQE vectors (Qiagen).

A variety of fungal cell expression vectors may be used to express recombinant human VR3 receptor in fungal cells such as yeast. Commercially available fungal cell expression vectors which may be suitable for recombinant human VR3 receptor expression include but are not limited to pYES2 (InVitrogen) and Pichia expression vector (InVitrogen). A variety of insect cell expression vectors may be used to express recombinant human VR3 receptor in insect cells. Commercially available insect cell expression vectors which may be suitable for recombinant expression of human VR3 receptor include but are not limited to pBlueBacII (InVitrogen).

DNA encoding human VR3 receptor may be cloned into an expression vector for expression in a recombinant host cell. Recombinant host cells may be prokaryotic or eukaryotic, including but not limited to bacteria such as E. coli, fungal cells such as yeast, mammalian cells including but not limited to cell lines of human, bovine, porcine, monkey and rodent origin, and insect cells including but not limited to drosophila and silkworm derived cell line. Cell lines derived from mammalian species which may be suitable and which are commercially available, include but are not limited to, CV-1 (ATCC CCL 70), COS-1 (ATCC CRL 1650), COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCC CCL 92), NIH/3T3 (ATCC CRL 1658), HeLa (ATCC CCL 2), C127I (ATCC CRL 1616), BS-C-1 (ATCC CCL 26), MRC-5 (ATCC CCL 171), L-cells, and HEK-293 (ATCC CRL1573).

The expression vector may be introduced into host cells via any one of a number of techniques including but not limited to transformation, transfection, protoplast fusion, lipofection, and electroporation. The expression vector-containing cells are clonally propagated and individually analyzed to determine whether they produce human VR3 receptor protein. Identification of human VR3 receptor expressing host cell clones may be done by several means, including but not limited

to immunological reactivity with anti-human VR3 receptor antibodies, and the presence of host cell-associated human VR3 receptor activity.

Expression of human VR3 receptor DNA may also be performed using *in vitro* produced synthetic mRNA. Synthetic mRNA or mRNA isolated from Human VR3 receptor producing cells can be efficiently translated in various cell-free systems, including but not limited to wheat germ extracts and reticulocyte extracts, as well as efficiently translated in cell based systems, including but not limited to microinjection into frog oocytes, with microinjection into frog oocytes being generally preferred.

To determine the human VR3 receptor DNA sequence(s) that yields optimal levels of human VR3 receptor activity and/or human VR3 receptor protein, human VR3 receptor DNA molecules including, but not limited to, the following can be constructed:

Gene name	Start codon	End codon	total base pairs
VR3A+B+	837	3065	2229
VR3A+B-	338	2953	2616
VR3A-B-	1	2436	2436

(these numbers correspond to first nucleotide of first methionine and last nucleotide before the first stop codon) and several constructs containing portions of the cDNA encoding human VR3 receptor protein. All constructs can be designed to contain none, all or portions of the 5' or the 3' untranslated region of human VR3 receptor cDNA. Human VR3 receptor activity and levels of protein expression can be determined following the introduction, both singly and in combination, of these constructs into appropriate host cells. Following determination of the human VR3 receptor DNA cassette yielding optimal expression in transient assays, this Human VR3 receptor DNA construct is transferred to a variety of expression vectors, for expression in host cells including, but not limited to, mammalian cells, baculovirus-infected insect cells, E. coli, and the yeast S. cerevisiae.

Assay methods for human VR3 receptor

Host cell transfectants and microinjected oocytes may be used to assay both the levels of functional Human VR3 receptor activity and levels of total human VR3 receptor protein by the following methods. In the case of recombinant host cells, this involves the co-transfection of one or possibly two or more plasmids, containing the human VR3 receptor DNA encoding one or more fragments, subunits, or other functional gene. In the case of oocytes, this involves the co-injection of synthetic RNAs for human VR3 receptor protein. Following an appropriate period of time to allow for expression, cellular protein is metabolically labelled with, for example ³⁵S-methionine for 24 hours, after which cell lysates and cell culture supernatants are harvested and subjected to immunoprecipitation with polyclonal antibodies directed against the human VR3 receptor protein.

Levels of human VR3 receptor protein in host cells are quantitated by immunoaffinity and/or ligand affinity techniques. Human VR3 receptor-specific affinity beads or human VR3 receptor-specific antibodies are used to isolate for example ³⁵S-methionine labelled or unlabelled human VR3 receptor protein. Labelled human VR3 receptor protein is analyzed by SDS-PAGE. Unlabelled human VR3 receptor protein is detected by Western blotting, ELISA or RIA assays employing human VR3 receptor specific antibodies.

Other methods for detecting human VR3 receptor activity involve the direct measurement of human VR3 receptor activity in whole cells transfected with human VR3 receptor cDNA or oocytes injected with human VR3 receptor mRNA. Human VR3 receptor activity is measured by specific ligand binding or biological characteristics of the host cells expressing human VR3 receptor DNA. In the case of recombinant host cells expressing human VR3 receptor patch voltage clamp techniques can be used to measure channel activity and quantitate human VR3 receptor protein. In the case of oocytes patch clamp as well as two-electrode voltage

clamp techniques can be used to measure calcium channel activity and quantitate human VR3 receptor protein.

Cell based assays

The present invention provides a whole cell method to detect compound modulation of human VR3 receptor. The method comprises the steps;

- 1) contacting a compound, and a cell that contains functional human VR3 receptor, and
- 2) measuring a change in the cell in response to modified human VR3 receptor function by the compound.

The amount of time necessary for cellular contact with the compound is empirically determined, for example, by running a time course with a known human VR3 receptor modulator and measuring cellular changes as a function of time.

The measurement means of the method of the present invention can be further defined by comparing a cell that has been exposed to a compound to an identical cell that has not been similarly exposed to the compound. Alternatively two cells, one containing functional human VR3 receptor and a second cell identical to the first, but lacking functional human VR3 receptor could be both be contacted with the same compound and compared for differences between the two cells. This technique is also useful in establishing the background noise of these assays. One of average skill in the art will appreciate that these control mechanisms also allow easy selection of cellular changes that are responsive to modulation of functional human VR3 receptor.

The term "cell" refers to at least one cell, but includes a plurality of cells appropriate for the sensitivity of the detection method. Cells suitable for the present invention may be bacterial, yeast, or eukaryotic.

The assay methods to determine compound modulation of functional human VR3 receptor can be in conventional laboratory format or adapted for high throughput. The term "high throughput" refers to an assay design that allows easy

analysis of multiple samples simultaneously, and capacity for robotic manipulation. Another desired feature of high throughput assays is an assay design that is optimized to reduce reagent usage, or minimize the number of manipulations in order to achieve the analysis desired. Examples of assay formats include but are not limited to, 96-well or 384-well plates, levitating droplets, and “lab on a chip” microchannel chips used for liquid handling experiments. It is well known by those in the art that as miniaturization of plastic molds and liquid handling devices are advanced, or as improved assay devices are designed, that greater numbers of samples may be performed using the design of the present invention.

The cellular changes suitable for the method of the present invention comprise directly measuring changes in the activity, function or quantity of human VR3 receptor, or by measuring downstream effects of human VR3 receptor function, for example by measuring secondary messenger concentrations or changes in transcription or by changes in protein levels of genes that are transcriptionally influenced by human VR3 receptor, or by measuring phenotypic changes in the cell. Preferred measurement means include changes in the quantity of human VR3 receptor protein, changes in the functional activity of human VR3 receptor, changes in the quantity of mRNA, changes in intracellular protein, changes in cell surface protein, or secreted protein, or changes in Ca^{+2} , cAMP or GTP concentration. Changes in the quantity or functional activity of human VR3 receptor are described herein. Changes in the levels of mRNA are detected by reverse transcription polymerase chain reaction (RT-PCR) or by differential gene expression. Immunoaffinity, ligand affinity, or enzymatic measurement quantitates VR3 induced changes in levels of specific proteins in host cells. Where the protein is an enzyme, the induction of protein is monitored by cleavage of a fluorogenic or colorimetric substrate.

Preferred detection means for cell surface protein include flow cytometry or statistical cell imaging. In both techniques the protein of interest is localized at the

cell surface, labeled with a specific fluorescent probe, and detected via the degree of cellular fluorescence. In flow cytometry, the cells are analyzed in a solution, whereas in cellular imaging techniques, a field of cells is compared for relative fluorescence.

The present invention is also directed to methods for screening for compounds that modulate the expression of DNA or RNA encoding human VR3 receptor as well as the function of human VR3 receptor protein *in vivo*.

Compounds may modulate by increasing or attenuating the expression of DNA or RNA encoding human VR3 receptor, or the function of human VR3 receptor protein. Compounds that modulate the expression of DNA or RNA encoding human VR3 receptor or the function of human VR3 receptor protein may be detected by a variety of assays. The assay may be a simple "yes/no" assay to determine whether there is a change in expression or function. The assay may be made quantitative by comparing the expression or function of a test sample with the levels of expression or function in a standard sample. Modulators identified in this process are useful as therapeutic agents, and human VR3 receptor.

Purification of human VR3 receptor protein

Following expression of human VR3 receptor in a recombinant host cell, human VR3 receptor protein may be recovered to provide purified human VR3 receptor. Recombinant human VR3 receptor may be purified from cell lysates and extracts, by various combinations of, or individual application of salt fractionation, ion exchange chromatography, size exclusion chromatography, hydroxylapatite adsorption chromatography and hydrophobic interaction chromatography, lectin chromatography, and antibody/ligand affinity chromatography.

Recombinant human VR3 receptor can be separated from other cellular proteins by use of an immunoaffinity column made with monoclonal or polyclonal antibodies specific for full length nascent human VR3 receptor, polypeptide fragments of human VR3 receptor or human VR3 receptor subunits. The affinity

resin is then equilibrated in a suitable buffer, for example phosphate buffered saline (pH 7.3), and the cell culture supernatants or cell extracts containing human VR3 receptor or human VR3 receptor subunits are slowly passed through the column. The column is then washed with the buffer until the optical density (A_{280}) falls to background, then the protein is eluted by changing the buffer condition, such as by lowering the pH using a buffer such as 0.23 M glycine-HCl (pH 2.6). The purified Human VR3 receptor protein is then dialyzed against a suitable buffer such as phosphate buffered saline.

Production and use of antibodies that bind to human VR3 receptor

Monospecific antibodies to human VR3 receptor are purified from mammalian antisera containing antibodies reactive against human VR3 receptor or are prepared as monoclonal antibodies reactive with human VR3 receptor using the technique originally described by Kohler and Milstein, *Nature* 256: 495-497 (1975). Immunological techniques are well known in the art and described in, for example, Antibodies: A laboratory manual published by Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, ISBN 0879693142. Monospecific antibody as used herein is defined as a single antibody species or multiple antibody species with homogenous binding characteristics for human VR3 receptor. Homogenous binding as used herein refers to the ability of the antibody species to bind to a specific antigen or epitope, such as those associated with the human VR3 receptor, as described above. Human VR3 receptor specific antibodies are raised by immunizing animals such as mice, rats, guinea pigs, rabbits, goats, horses and the like, with rabbits being preferred, with an appropriate concentration of human VR3 receptor either with or without an immune adjuvant.

Preimmune serum is collected prior to the first immunization. Each animal receives between about 0.001 mg and about 100 mg of human VR3 receptor

associated with an acceptable immune adjuvant. Such acceptable adjuvants include, but are not limited to, Freund's complete, Freund's incomplete, alum-precipitate, water in oil emulsion containing Corynebacterium parvum and tRNA. The initial immunization consists of human VR3 receptor in, preferably, Freund's complete adjuvant at multiple sites either subcutaneously (SC), intraperitoneally (IP) or both. Each animal is bled at regular intervals, preferably weekly, to determine antibody titer. The animals may or may not receive booster injections following the initial immunization. Those animals receiving booster injections are generally given an equal amount of the antigen in Freund's incomplete adjuvant by the same route. Booster injections are given at about three-week intervals until maximal titers are obtained. At about 7 days after each booster immunization or about weekly after a single immunization, the animals are bled, the serum collected, and aliquots are stored at about -20°C.

Monoclonal antibodies (mAb) reactive with human VR3 receptor are prepared by immunizing inbred mice, preferably Balb/c, with human VR3 receptor. The mice are immunized by the IP or SC route with about 0.001 mg to about 1.0 mg, preferably about 0.1 mg, of human VR3 receptor in about 0.1 ml buffer or saline incorporated in an equal volume of an acceptable adjuvant, as discussed above. Freund's adjuvant is preferred, with Freund's complete adjuvant being used for the initial immunization and Freund's incomplete adjuvant used thereafter. The mice receive an initial immunization on day 0 and are rested for about 2 to about 30 weeks. Immunized mice are given one or more booster immunizations of about 0.001 to about 1.0 mg of human VR3 receptor in a buffer solution such as phosphate buffered saline by the intravenous (IV) route. Lymphocytes, from antibody positive mice, preferably splenic lymphocytes, are obtained by removing spleens from immunized mice by standard procedures known in the art. Hybridoma cells are produced by mixing the splenic lymphocytes with an appropriate fusion partner, preferably myeloma cells, under conditions that will allow the formation of stable hybridomas. Fusion partners

may include, but are not limited to: mouse myelomas P3/NS1/Ag 4-1; MPC-11; S-194 and Sp2/0, with Sp2/0 being generally preferred. The antibody producing cells and myeloma cells are fused in polyethylene glycol, about 1000 mol. wt., at concentrations from about 30% to about 50%. Fused hybridoma cells are selected by growth in hypoxanthine, thymidine and aminopterin supplemented Dulbecco's Modified Eagles Medium (DMEM) by procedures known in the art. Supernatant fluids are collected from growth positive wells on about days 14, 18, and 21 and are screened for antibody production by an immunoassay such as solid phase immunoradioassay (SPIRA) using human VR3 receptor as the antigen. The culture fluids are also tested in the Ouchterlony precipitation assay to determine the isotype of the mAb. Hybridoma cells from antibody positive wells are cloned by a technique such as the soft agar technique of MacPherson, Soft Agar Techniques, in Tissue Culture Methods and Applications, Kruse and Paterson, Eds., Academic Press, 1973 or by the technique of limited dilution.

Monoclonal antibodies are produced *in vivo* by injection of pristane primed Balb/c mice, approximately 0.5 ml per mouse, with about 1×10^6 to about 6×10^6 hybridoma cells at least about 4 days after priming. Ascites fluid is collected at approximately 8-12 days after cell transfer and the monoclonal antibodies are purified by techniques known in the art.

In vitro production of anti-human VR3 receptor mAb is carried out by growing the hybridoma in tissue culture media well known in the art. High density *in vitro* cell culture may be conducted to produce large quantities of anti-human VR3 receptor MAbs using hollow fiber culture techniques, air lift reactors, roller bottle, or spinner flasks culture techniques well known in the art. The mAb are purified by techniques known in the art.

Antibody titers of ascites or hybridoma culture fluids are determined by various serological or immunological assays which include, but are not limited to, precipitation, passive agglutination, enzyme-linked immunosorbent antibody (ELISA)

technique and radioimmunoassay (RIA) techniques. Similar assays are used to detect the presence of human VR3 receptor in body fluids or tissue and cell extracts.

It is readily apparent to those skilled in the art that the above described methods for producing monospecific antibodies may be utilized to produce antibodies specific for human VR3 receptor polypeptide fragments, or full-length nascent human VR3 receptor polypeptide, or the individual human VR3 receptor subunits. Specifically, it is readily apparent to those skilled in the art that monospecific antibodies may be generated which are specific for only one human VR3 receptor subunit or the fully functional human VR3 receptor protein. It is also apparent to those skilled in the art that monospecific antibodies may be generated that inhibit normal function of human VR3 receptor protein.

Human VR3 receptor antibody affinity columns are made by adding the antibodies to a gel support such that the antibodies form covalent linkages with the gel bead support. Preferred covalent linkages are made through amine, aldehyde, or sulfhydryl residues contained on the antibody. Methods to generate aldehydes or free sulfhydryl groups on antibodies are well known in the art; amine groups are reactive with, for example, N-hydroxysuccinimide esters.

Kit compositions containing human VR3 receptor specific reagents

Kits containing human VR3 receptor DNA or RNA, antibodies to human VR3 receptor, or human VR3 receptor protein may be prepared. Such kits are used to detect DNA that hybridizes to human VR3 receptor DNA or to detect the presence of human VR3 receptor protein or peptide fragments in a sample. Such characterization is useful for a variety of purposes including but not limited to forensic analyses, diagnostic applications, and epidemiological studies.

This invention relates to the use of human VR3 polynucleotides for the use as diagnostic reagents. Detection of a mutated form of human VR3 gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a

disease or susceptibility to a disease which results from under-expression or over-expression of human VR3. Individuals carrying mutations in the human VR3 gene may be detected at the DNA level by a variety of techniques well known in the art, and described herein.

The DNA molecules, RNA molecules, recombinant protein and antibodies of the present invention may be used to screen and measure levels of human VR3 receptor DNA, human VR3 receptor RNA or human VR3 receptor protein. The recombinant proteins, DNA molecules, RNA molecules and antibodies lend themselves to the formulation of kits suitable for the detection and typing of human VR3 receptor. Such a kit would comprise a compartmentalized carrier suitable to hold in close confinement at least one container. The carrier would further comprise reagents such as recombinant human VR3 receptor protein or anti-human VR3 receptor antibodies suitable for detecting human VR3 receptor. The carrier may also contain a means for detection such as labeled antigen or enzyme substrates or the like.

Gene therapy

Nucleotide sequences that are complementary to the human VR3 receptor encoding DNA sequence can be synthesized for antisense therapy. These antisense molecules may be DNA, stable derivatives of DNA such as phosphorothioates or methylphosphonates, RNA, stable derivatives of RNA such as 2'-O-alkylRNA, or other Human VR3 receptor antisense oligonucleotide mimetics. Human VR3 receptor antisense molecules may be introduced into cells by microinjection, liposome encapsulation or by expression from vectors harboring the antisense sequence. Human VR3 receptor antisense therapy may be particularly useful for the treatment of diseases where it is beneficial to reduce human VR3 receptor activity.

Human VR3 receptor gene therapy may be used to introduce human VR3 receptor into the cells of target organisms. The human VR3 receptor gene can be ligated into viral vectors that mediate transfer of the human VR3 receptor DNA by

infection of recipient host cells. Suitable viral vectors include retrovirus, adenovirus, adeno-associated virus, herpes virus, vaccinia virus, poliovirus and the like. Alternatively, human VR3 receptor DNA can be transferred into cells for gene therapy by non-viral techniques including receptor-mediated targeted DNA transfer using ligand-DNA conjugates or adenovirus-ligand-DNA conjugates, lipofection membrane fusion or direct microinjection. These procedures and variations thereof are suitable for *ex vivo* as well as *in vivo* human VR3 receptor gene therapy. Human VR3 receptor gene therapy may be particularly useful for the treatment of diseases where it is beneficial to elevate human VR3 receptor activity. Protocols for molecular methodology of gene therapy suitable for use with the human VR3 receptor gene is described in Gene Therapy Protocols, edited by Paul D. Robbins, Human press, Totawa NJ, 1996.

Pharmaceutical compositions

Pharmaceutically useful compositions comprising human VR3 receptor DNA, human VR3 receptor RNA, or human VR3 receptor protein, or modulators of human VR3 receptor activity, may be formulated according to known methods such as by the admixture of a pharmaceutically acceptable carrier. Examples of such carriers and methods of formulation may be found in Remington's Pharmaceutical Sciences. To form a pharmaceutically acceptable composition suitable for effective administration, such compositions will contain an effective amount of the protein, DNA, RNA, or modulator.

Therapeutic or diagnostic compositions of the invention are administered to an individual in amounts sufficient to treat or diagnose disorders in which modulation of Human VR3 receptor-related activity is indicated. The effective amount may vary according to a variety of factors such as the individual's condition, weight, sex and age. Other factors include the mode of administration. The pharmaceutical

compositions may be provided to the individual by a variety of routes such as subcutaneous, topical, oral and intramuscular.

The term "chemical derivative" describes a molecule that contains additional chemical moieties that are not normally a part of the base molecule. Such moieties may improve the solubility, half-life, absorption, etc. of the base molecule. Alternatively the moieties may attenuate undesirable side effects of the base molecule or decrease the toxicity of the base molecule. Examples of such moieties are described in a variety of texts, such as Remington's Pharmaceutical Sciences.

Compounds identified according to the methods disclosed herein may be used alone at appropriate dosages defined by routine testing in order to obtain optimal inhibition of the human VR3 receptor or its activity while minimizing any potential toxicity. In addition, co-administration or sequential administration of other agents may be desirable.

The present invention also has the objective of providing suitable topical, oral, systemic and parenteral pharmaceutical formulations for use in the novel methods of treatment of the present invention. The compositions containing compounds or modulators identified according to this invention as the active ingredient for use in the modulation of human VR3 receptor can be administered in a wide variety of therapeutic dosage forms in conventional vehicles for administration. For example, the compounds or modulators can be administered in such oral dosage forms as tablets, capsules (each including timed release and sustained release formulations), pills, powders, granules, elixirs, tinctures, solutions, suspensions, syrups and emulsions, or by injection. Likewise, they may also be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous, topical with or without occlusion, or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the compound desired can be employed as a human VR3 receptor-modulating agent.

The daily dosage of the products may be varied over a wide range from 0.01 to 1,000 mg per patient, per day. For oral administration, the compositions are preferably provided in the form of scored or unscored tablets containing 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, and 50.0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.0001 mg/kg to about 100 mg/kg of body weight per day. The range is more particularly from about 0.001 mg/kg to 10 mg/kg of body weight per day. The dosages of the human VR3 receptor modulators are adjusted when combined to achieve desired effects. On the other hand, dosages of these various agents may be independently optimized and combined to achieve a synergistic result wherein the pathology is reduced more than it would be if either agent were used alone.

Advantageously, compounds or modulators of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds or modulators for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

For combination treatment with more than one active agent, where the active agents are in separate dosage formulations, the active agents can be administered concurrently, or they each can be administered at separately staggered times.

The dosage regimen utilizing the compounds or modulators of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound thereof employed. A physician or veterinarian of

ordinary skill can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentrations of drug within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a drug.

In the methods of the present invention, the compounds or modulators herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include, without limitation, sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

For liquid forms the active drug component can be combined in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. Other dispersing agents that may be employed include glycerin and the like. For parenteral administration, sterile suspensions

and solutions are desired. Isotonic preparations, which generally contain suitable preservatives, are employed when intravenous administration is desired.

Topical preparations containing the active drug component can be admixed with a variety of carrier materials well known in the art, such as, e.g., alcohols, aloe vera gel, allantoin, glycerine, vitamin A and E oils, mineral oil, PPG2 myristyl propionate, and the like, to form, e.g., alcoholic solutions, topical cleansers, cleansing creams, skin gels, skin lotions, and shampoos in cream or gel formulations.

The compounds or modulators of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds or modulators of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinyl-pyrrolidone, pyran copolymer, polyhydroxypropylmethacryl-amidephenol, polyhydroxy-ethylaspartamidephenol, or polyethyl-eneoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds or modulators of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydro-pyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

For oral administration, the compounds or modulators may be administered in capsule, tablet, or bolus form or alternatively they can be mixed in the animals feed. The capsules, tablets, and boluses are comprised of the active ingredient in combination with an appropriate carrier vehicle such as starch, talc, magnesium stearate, or di-calcium phosphate. These unit dosage forms are

prepared by intimately mixing the active ingredient with suitable finely-powdered inert ingredients including diluents, fillers, disintegrating agents, and/or binders such that a uniform mixture is obtained. An inert ingredient is one that will not react with the compounds or modulators and which is non-toxic to the animal being treated. Suitable inert ingredients include starch, lactose, talc, magnesium stearate, vegetable gums and oils, and the like. These formulations may contain a widely variable amount of the active and inactive ingredients depending on numerous factors such as the size and type of the animal species to be treated and the type and severity of the infection. The active ingredient may also be administered as an additive to the feed by simply mixing the compound with the feedstuff or by applying the compound to the surface of the feed. Alternatively the active ingredient may be mixed with an inert carrier and the resulting composition may then either be mixed with the feed or fed directly to the animal. Suitable inert carriers include corn meal, citrus meal, fermentation residues, soya grits, dried grains and the like. The active ingredients are intimately mixed with these inert carriers by grinding, stirring, milling, or tumbling such that the final composition contains from 0.001 to 5% by weight of the active ingredient.

The compounds or modulators may alternatively be administered parenterally via injection of a formulation consisting of the active ingredient dissolved in an inert liquid carrier. Injection may be either intramuscular, intraruminal, intratracheal, or subcutaneous. The injectable formulation consists of the active ingredient mixed with an appropriate inert liquid carrier. Acceptable liquid carriers include the vegetable oils such as peanut oil, cottonseed oil, sesame oil and the like as well as organic solvents such as solketal, glycerol formal and the like. As an alternative, aqueous parenteral formulations may also be used. The vegetable oils are the preferred liquid carriers. The formulations are prepared by dissolving or suspending the active ingredient in the liquid carrier such that the final formulation contains from 0.005 to 10% by weight of the active ingredient.

Topical application of the compounds or modulators is possible through the use of a liquid drench or a shampoo containing the instant compounds or modulators as an aqueous solution or suspension. These formulations generally contain a suspending agent such as bentonite and normally will also contain an antifoaming agent. Formulations containing from 0.005 to 10% by weight of the active ingredient are acceptable. Preferred formulations are those containing from 0.01 to 5% by weight of the instant compounds or modulators.

The following examples illustrate the present invention without, however, limiting the same thereto.

EXAMPLE 1 Generation of human prostate and pituitary cDNA libraries

cDNA synthesis:

First strand synthesis: Approximately 5 µg of human prostate or pituitary mRNA (Clontech) was used to synthesize cDNA using the cDNA synthesis kit (Life Technologies). Two microliters of NotI primer adapter was added to 5µl of mRNA and the mixture was heated to 70 °C for 10 minutes and placed on ice. The following reagents were added on ice: 4µl of 5x first strand buffer (250mM TRIS-HCl (pH8.3), 375mM KCl, 15mM MgCl₂), 2µl of 0.1M DTT, 10mM dNTP (nucleotide triphosphates) mix and 1µl of DEPC treated water. The reaction was incubated at 42 °C for 5minutes. Finally, 5µl of Superscript RT II was added and incubated at 42 °C for 2 more hours. The reaction was terminated on ice.

Second strand synthesis: The first strand product was adjusted to 93µl with water and the following reagents were added on ice: 30 µl of 5x 2nd strand buffer (100 mM TRIS-HCl (pH6.9), 450 mM KCl, 23 mM MgCl₂, 0.75 mM β-NAD⁺, 50mM (NH₄)₂SO₄), 3µl of 10 mM dNTP (nucleotide triphosphates), 1µl *E. coli* DNA ligase (10units) 1µl RNase H (2units), 4 µl DNA pol I (10 units). The reaction was

incubated at 16°C for 2 hours. The DNA from second strand synthesis was treated with T4 DNA polymerase and placed at 16°C to blunt the DNA ends. The double stranded cDNA was extracted with 150 µl of a mixture of phenol and chloroform (1:1, v:v) and precipitated with 0.5 volumes of 7.5 M NH₄OAc and 2 volumes of absolute ethanol. The pellet was washed with 70% ethanol and dried down at 37°C to remove the residual ethanol. The double stranded DNA pellet was resuspended in 25 µl of water and the following reagents were added; 10 µl of 5x T4 DNA ligase buffer, 10 µl of SalI adapters and 5 µl of T4 DNA ligase. The ingredients were mixed gently and ligated overnight at 16° C. The ligation mix was extracted with phenol:chloroform:isoamyl alcohol, vortexed thoroughly and centrifuged at room temperature for 5 minutes at 14,000 x g to separate the phases. The aqueous phase was transferred to a new tube and the volume adjusted to 100 µl with water. The purified DNA was size selected on a chromaspin 1000 column (Clontech) to eliminate the smaller cDNA molecules. The double stranded DNA was digested with NotI restriction enzyme for 3-4 hours at 37° C. The restriction digest was electrophoresed on a 0.8 % low melt agarose gel. The cDNA in the range of 1-5 kb was cut out and purified using Gelzyme (In Vitrogen). The product was extracted with phenol:chloroform and precipitated with NH₄OAc and absolute ethanol. The pellet was washed with 70% ethanol and resuspended in 10 µl of water.

Ligation of cDNA to the Vector: The cDNA was split up into 5 tubes (2µl each) and the ligation reactions were set up by adding 4.5 µl of water, 2 µl of 5x ligation buffer, 1µl of p-Sport vector DNA (cut with Sal-I / NotI and phosphatase treated) and 0.5 µl of T4 DNA ligase. The ligation was incubated at 40° C overnight.

Introduction of Ligated cDNA into E. coli by Electroporation:

The ligation reaction volume was adjusted to a total volume of 20 µl with water. Five milliliters of yeast tRNA, 12.5 ml of 7.5M NH₄OAc and 70 ml of absolute ethanol (-20°C) was added. The mixture was vortexed thoroughly, and immediately centrifuged at room temperature for 20 minutes at 14000 xg. The pellets were washed in 70% ethanol and each pellet was resuspended in 5 ml of water. All 5 ligations (25ml) were pooled and 100µl of DH10B electro-competent cells (Life Technologies) were electroporated with 1 ml of DNA (total of 20 electroporations), then plated out on ampicillin plates to determine the number of recombinants (cfu) per microliter. The entire library was seeded into 2 liters of Super Broth and maxipreps were made using Promega Maxi Prep kit and purified on cesium chloride gradients.

EXAMPLE 2: Library Screening / human VR3 A+B+ Generation

Human pituitary gland library screening:

One-microliter aliquots of the human pituitary gland library were electroporated into Electromax DH10B cells (Life Technologies). The volume was adjusted to 1 ml with SOC media and incubated for 45 minutes at 37°C with shaking. The library was then plated out on 150cm² plates containing LB to a density of 20000 colonies per plate. These cultures were grown overnight at 37°C.

A human VR3 receptor probe was generated by polymerase chain reaction using the following primer pair:

5' oligo (SEQ.ID. NO.:1): 5' ACCGGCCTATCCTCTTTGACATCGTG

3' oligo (SEQ.ID.NO.:2): 5' TGTCCGCCTTCTTGTGGGGGTTCTC

The probe was generated by PCR using regular PCR conditions using 5' and 3' probe oligos (100ng each) and 10 ng of diluted miniprep DNA. The resulting 493 bp fragment was run on 1% agarose gel and purified using a QIAquick Gel extraction kit (Quiagen). About 100 ng of the purified probe was labeled with alpha 32P using

oligolabeling kit from Pharmacia and the labeled DNA was purified with S-200 columns (Pharmacia).

The library colonies were lifted on Protran nitrocellulose filters (Scheicher & Schuel) and the DNA was denatured in 1.5 M NaCl, 0.5 M NaOH. The filter disks were neutralized with 1.5 M NaCl, 1.0 M Tris-HCl, pH 7.5 and then UV crosslinked to the membrane using a UV-Stratalinker (Stratagene). The filters were washed several times in wash solution (1 M Tris-HCl, pH 8.0; 5 M NaCl; 0.5 M EDTA; 20% SDS) at 42°C. Then the disks were incubated in 1x southern pre-hybridization buffer (5'-3' Inc) containing 50% formamide and 100 ug/ml of sheared salmon sperm DNA (5' - 3' Inc) for 6 hours at 42 C. Finally, hybridization was performed overnight at 42C in 1x hybridization buffer (5'-3') containing 50% formamide, 100ng of sheared salmon sperm DNA in the presence of labeled probe (5×10^5 to 1×10^6 cmp/ml of hybridization buffer).

The disks were washed twice in 2xSSC, 0.2% SDS at room temperature (20 minutes each) and once in 0.2xSSC, 0.1%SDS at 50C for 30 minutes. The membranes were than placed on sheets of filter paper, wrapped in the plastic wrap and exposed to the film at -20C overnight.

Positive clones were identified and collected by coring the colonies from the original plate. The colonies were incubated in LB for 1 hour at 37°C. Dilutions of the cultures were plated onto LB agar plates and the filter-lifting, hybridizing, washing, colony-picking procedure was repeated. Individual clones from the second screen were picked and digested with SalI/NotI to determine the size of the inserts, and the inserts were sequenced.

The full length clone was generated by PCR with Pfu polymerase using 10 ng of the sequenced library clone as a template and full length oligos with KpnI (FL 5' oligo SEQ.ID.NO.3) and NotI (FL 3' oligo SEQ.ID.NO.4) sites.

FL 5' oligo (SEQ.ID.NO.3):

AACGTTGGTACCGCCACCATGGCGGATTCCAGCGAAGGCCCCCGCGCG

FL3' oligo: (SEQ.ID.NO.4):

TAAAGCGGCCGCTTCAGGAGGGACATCGGTGAGCCTCAC

The PCR product was digested with KpnI and NotI enzymes and cloned into a pSP64T.GC expression vector. Large-scale preparation of DNA was done using a MEGA prep kit (Quiagen).

EXAMPLE 3: Library Screening / human VR3 A+B- and human VR3 A-B- Generation
Human prostate library screening:

One microliter aliquots of the human prostate library were electroporated into Electromax DH10B cells (Life Technologies). The volume was adjusted to 1 ml with SOC media and incubated for 45 minutes at 37°C with shaking. The library was then plated out on 150cm² plates containing LB to a density of 20000 colonies per plate. These cultures were grown overnight at 37°C.

A human VR3 receptor probe was generated by polymerase chain reaction using the following primer pair:

5' oligo (SEQ.ID. NO.:13): 5' CTACCTGACGGAGAACCCCCACAAG

3' oligo (SEQ.ID.NO.:14): 5' GTAGTAGGCGGTGAGACTGAAGATGA

The probe was generated by PCR using regular PCR conditions using 5' and 3' probe oligos (100ng each) and 10 ng of diluted miniprep DNA. The resulting 387 bp fragment was run on 1% agarose gel and purified using a QUIAquick Gel extraction kit (Quiagen). About 100 ng of the purified probe was labeled with alpha 32P using oligolabeling kit from Pharmacia and the labeled DNA was purified with S-200 columns (Pharmacia).

The library colonies were lifted on Protran nitrocellulose filters (Schleicher & Schuel) and the DNA was denatured in 1.5 M NaCl, 0.5 M NaOH. The filter disks were neutralized with 1.5 M NaCl, 1.0 M Tris-HCl, pH 7.5 and then UV crosslinked to the membrane using a UV-Stratalinker (Stratagene). The filters were washed several times in wash solution (1 M Tris-HCl, pH 8.0; 5 M NaCl; 0.5 M EDTA; 20% SDS) at 42°C. Then the disks were incubated in 1x southern pre-hybridization buffer (5'-3' Inc) containing 50% formamide and 100 ug/ml of sheared salmon sperm DNA (5' - 3' Inc) for 6 hours at 42 C. Finally, hybridization was performed overnight at 42C in 1x hybridization buffer (5'-3') containing 50% formamide, 100ng of sheared salmon sperm DNA in the presence of labeled probe (5×10^5 to 1×10^6 cmp/ml of hybridization buffer).

The disks were washed twice in 2xSSC, 0.2% SDS at room temperature (20 minutes each) and once in 0.2xSSC, 0.1%SDS at 50C for 30 minutes. The membranes were than placed on sheets of filter paper, wrapped in the Saran wrap and exposed to the film at -20C overnight.

Positive clones were identified and collected by coring the colonies from the original plate. The colonies were incubated in LB for 1 hour at 37°C. Dilutions of the cultures were plated onto LB agar plates and the filter-lifting, hybridizing, washing, colony-picking procedure was repeated. Individual clones from the second screen were picked and digested with EcoRI/NotI to determine the size of the inserts, and the inserts were sequenced.

The full length clone was generated by PCR with Pfu polymerase using 10 ng of the sequenced library clone as a template and full length oligos with NotI (FL 5' oligo SEQ.ID.NO.:15) and XbaI (FL 3' oligo SEQ.ID.NO.:16) sites.

FL 5' oligo (SEQ.ID.NO.:15):

5' AACGTTGGCGGCCGCGCCACCATGGCGGATTCCAGCGAAGGCCCCCGCGC
G

FL3' oligo: (SEQ.ID.NO.:16):

5' AACGTTTCTAGACTGGGCTGCAGTCCCTAG

The PCR product was digested with NotI and XbaI enzymes and cloned into a pGem HE expression vector. Large-scale preparation of DNA was done using a MEGA prep kit (Quiagen).

EXAMPLE 4- Cloning human VR3 receptor cDNA into a Mammalian Expression Vector

The human VR3 receptor cDNAs (collectively referred to as hVR3) were cloned into the mammalian expression vector pcDNA3.1/Zeo(+). The cloned PCR product was purified on a column (Wizard PCR DNA purification kit from Promega) and digested with Not I and EcoRI (NEB) to create cohesive ends. The product was purified by electrophoresis on a low melting point agarose gel. The pcDNA3.1/Zeo(+) vector was digested with NotI and XbaI (except for hVR3 A+B+, which was cloned into BamHI / NotI sites) enzymes and subsequently purified on a low melting point agarose gel. The linear vector was used to ligate to the human VR3 cDNA inserts. Recombinants were isolated, designated human VR3 receptor, and used to transfect mammalian cells (HEK293, COS-7 or CHO-K1 cells) using the Effectene non-liposomal lipid based transfection kit (Quiagen). Stable cell clones were selected by growth in the presence of zeocin. Single zeocin resistant clones were isolated and shown to contain the intact human VR3 receptor gene. Clones containing the human VR3 receptor cDNAs were analyzed for hVR3 protein expression. Recombinant plasmids containing human VR3 encoding DNA were used to transform the mammalian COS or CHO cells or HEK293 cells.

Cells expressing human VR3 receptor, stably or transiently, are used to test for expression of human VR3 receptor and for ³H-RTX binding activity. These cells are used to identify and examine other compounds for their ability to modulate,

inhibit or activate the human VR3 receptor and to compete for radioactive ^3H -RTX binding.

Cassettes containing the human VR3 receptor cDNA in the positive orientation with respect to the promoter are ligated into appropriate restriction sites 3' of the promoter and identified by restriction site mapping and/or sequencing. These cDNA expression vectors are introduced into fibroblastic host cells for example COS-7 (ATCC# CRL1651), and CV-1 tat [Sackevitz et al., Science 238: 1575 (1987)], 293, L (ATCC# CRL6362)] by standard methods including but not limited to electroporation, or chemical procedures (cationic liposomes, DEAE dextran, calcium phosphate). Transfected cells and cell culture supernatants are harvested and analyzed for human VR3 receptor expression as described herein.

All of the vectors used for mammalian transient expression can be used to establish stable cell lines expressing human VR3 receptor. Unaltered human VR3 receptor cDNA constructs cloned into expression vectors are expected to program host cells to make human VR3 receptor protein. The transfection host cells include, but are not limited to, CV-1-P [Sackevitz et al., Science 238: 1575 (1987)], tk-L [Wigler, et al. Cell 11: 223 (1977)], NS/0, and dHFr- CHO [Kaufman and Sharp, J. Mol. Biol. 159: 601, (1982)].

Co-transfection of any vector containing human VR3 receptor cDNA with a drug selection plasmid including, but not limited to G418, aminoglycoside phosphotransferase; hygromycin, hygromycin-B phosphotransferase; APRT, xanthine-guanine phosphoribosyl-transferase, will allow for the selection of stably transfected clones. Levels of human VR3 receptor are quantitated by the assays described herein.

Human VR3 receptor cDNA constructs are also ligated into vectors containing amplifiable drug-resistance markers for the production of mammalian cell clones synthesizing the highest possible levels of human VR3 receptor. Following introduction of these constructs into cells, clones containing the plasmid are selected

with the appropriate agent, and isolation of an over-expressing clone with a high copy number of plasmids is accomplished by selection in increasing doses of the agent.

The expression of recombinant human VR3 receptor is achieved by transfection of full-length human VR3 receptor cDNA into a mammalian host cell.

EXAMPLE 5- Characterization of functional protein encoded by pVR3R in *Xenopus* oocytes

Xenopus laevis oocytes were prepared and injected using standard methods previously described and known in the art (Fraser et al., 1993). Ovarian lobes from adult female *Xenopus laevis* (Nasco, Fort Atkinson, WI) were teased apart, rinsed several times in nominally Ca-free saline containing: 82.5mM NaCl, 2.5mM KCl, 1mM MgCl₂, 5 mM HEPES, adjusted to pH 7.0 with NaOH (OR-2), and gently shaken in OR-2 containing 0.2% collagenase Type 1 (ICN Biomedicals, Aurora, Ohio) for 2-5 hours. When approximately 50% of the follicular layers were removed, Stage V and VI oocytes were selected and rinsed in media consisting of 75% OR-2 and 25% ND-96. The ND-96 contained: 100 mM NaCl, 2 mM KCl, 1 mM MgCl₂,

1.8 mM CaCl₂, 5 mM HEPES, 2.5 mM Na pyruvate, gentamicin (50 ug/ml), adjusted to pH 7.0 with NaOH. The extracellular Ca⁺² was gradually increased and

1.8 mM CaCl₂, 5 mM HEPES, 2.5 mM Na pyruvate, gentamicin (50 ug/ml), adjusted to pH 7.0 with NaOH. The extracellular Ca⁺² was gradually increased and the cells were maintained in ND-96 for 2-24 hours before injection. For in vitro transcription, pGEM HE (Liman et al., 1992) containing human VR3 was linearized with NheI and transcribed with T7 RNA polymerase (Promega) in the presence of the cap analog m7G(5')ppp(5')G. The synthesized cRNA was precipitated with ammonium acetate and isopropanol, and resuspended in 50µl nuclease-free water. cRNA was quantified using formaldehyde gels (1% agarose, 1xMOPS, 3% formaldehyde) against 1.2 and 5 µl RNA markers (Gibco BRL, 0.24 - 9.5 Kb).

Oocytes were injected with 50 nl of the human VR3 receptor RNA (0.3 - 5 ng) with or without co-injection of VR1 (2-3 ng). Control oocytes were injected with 50 nl

of water. Oocytes were incubated for 1-13 days in ND-96 before analysis for expression of the human VR3. Incubations and collagenase digestion were carried out at room temperature. Injected oocytes were maintained in 48 well cell culture clusters (Costar; Cambridge, MA) at 18°C. Whole cell agonist-induced currents were measured 1-14 days after injection with a conventional two-electrode voltage clamp (GeneClamp500, Axon Instruments, Foster City, CA) using standard methods previously described and known in the art (Dascal, 1987). The microelectrodes were filled with 3 M KCl, which had resistances of 1 and 2 MΩ. Cells were continuously perfused with ND96 at ~10 ml/min at room temperature unless indicated. Membrane voltage was clamped at -70 mV unless indicated.

Oocytes were challenged with a variety of ligands, low pH, and depolarizing as well as hyperpolarizing voltage steps but there were no detectable differences in membrane conductance between human VR3-expressing oocytes and control oocytes [see Table 1]. However, human VR3 injected oocytes had properties that were different from controls in 4 respects. First, injection of VR3 A+B- cRNA caused oocytes to die. The viability of oocytes injected with VR3 isoforms having the A insert was significantly reduced compared to sister controls 3-4 days after injection (Figure 9). Second, all three isoforms of VR3 enhanced the heat-induced response (Figure 10). Third, A+B- co-injection with human VR1 produced a ruthenium red sensitive perfusion-induced current (lperfusion) (Figure 11). Fourth, human VR3 A+B- cRNA co-injection with human VR1 cRNA altered the responsiveness of the oocytes to 1 uM capsaicin, an agonist at the VR1 receptor (Figure 12).

TABLE 1

Stimulus	VR3 A+B- (3.25- 4 ng injected per oocyte unless specified)	VR3 A-B- (1.5 ng injected per oocyte)	VR3 A+B+ (5 ng injected per oocyte)
Capsaicin (1 uM)	NE: n=3 (n=3 0.4 ng)	NE: n=3	NE: n=3
Eugenol (10 uM)	NE: n=3 (n=3 0.4 ng)	NE: n=3	NE: n=3
Olvanil (10 uM)	NE: n=3 (n=1 0.4 ng)	NE: n=3	NE: n=3
Resiniferatoxin (1 uM)	NE: n=3 (n=1 0.4 ng)	NE: n=3	NE: n=3
Piperine (10 uM)	NE: n=3 (n=1 0.4 ng)	NE: n=3	NE: n=3
Gingerol (10 uM)	NE: n=3 (n=1 0.4 ng)	NE: n=3	NE: n=3
Guaiacol (10 uM)	NE: n=3 (n=1 0.4 ng)	NE: n=3	NE: n=3
β -phenylethylamine (100 uM)	NE: n=2	NT	NT
γ -hydroxybutyrate (100 uM)	NE: n=2	NT	NT
Anandamide (1.15 uM; RBI)	NE: n=3 (0.4 ng injected)	NE: n=1	NT

TABLE 1

Stimulus	VR3 A+B- (3.25- 4 ng injected per oocyte unless specified)	VR3 A-B- (1.5 ng injected per oocyte)	VR3 A+B+ (5 ng injected per oocyte)
Arachadonic acid (10 uM)	NT	NE: n=3	NT
pH 4.5 – 5.5	NE: n=2 (4 ng cRNA); n=1 (0.4 ng cRNA)	NE: n=1	NE: n=1
Depolarization	No difference from control (n=4)	No difference from control (n=6)	No difference from control (n=3)
Hyperpolarization	No difference from control (n=4)	No difference from control (n=6)	No difference from control (n=3)
NE: no effect; NT: not tested			

Viability.

Functional expression of human VR3 isoforms in *Xenopus* oocytes is shown: viability of oocytes maintained in ND-96 containing 2 Ca^{2+} was significantly diminished 4-6 days after injection with VR3 A+B- (3.25 ng) and VR3 A+B+ (5 ng) but not VR3 A-B- (1.5 ng) cRNA. Data for VR3 A+B- and control water-injected oocytes were similar in 5 separate experiments and combined. Dead oocytes were determined visually using a Bauch and Lomb dissecting microscope. Data were analyzed using Chi square analysis. VR3 A+B- injected oocytes were the least viable: only about 9% of oocytes survived to 4-6 days after injection. About 33% of oocytes injected with human VR3 A+B+ survived to 4-6 days after injection. About 67% of sister water-injected control oocytes survived the same time period under

essentially identical conditions. VR3 A-B- injected oocytes revealed similar viability to water injected controls. Similar data were obtained from 2 separate batches of cRNA.

Heat responsiveness.

Functional expression of human VR3 isoforms in *Xenopus* oocytes is shown: activation by heat. a). Shown is the mean peak current elicited by a heat ramp from 25 deg C to 46 deg C and maintained at 46 degC for at least 15 sec. Voltage ramps were applied from -120 to +80 mV over 400 msec at a sampling rate of 2 sec. The current shown was the increase in current over and above the initial current elicited at +80 mV (the rightmost point on the current voltage curve shown in (b)). Oocytes were injected with 3.25, 1.5 and 5 ng VR3 A+B-, A-B-, and A+B+ cRNA, respectively, and recorded up to 6 days later. Solid bar: water-injected controls (n=8); clear bar: VR3 A+B- isoform (n= 11); hatched bar: VR3 A-B- isoform (n= 8); stippled bar: VR3 A+B+ isoform (n=5). All isoforms showed a significant increase over water controls (p= 0.001, 0.03 and 0.007, respectively). The heat-induced response was about 2-fold larger in the A+B- isoform compared to the other 2 isoforms, but the differences were not significant (p= 0.051 and p= 0.16, for A+B- compared to A-B- and A+B+, respectively). Data were obtained from 2 sets of injected oocytes. Data shown is the mean and standard error of the mean. (b). Voltage-ramp induced currents were recorded during application of increasing heat to oocytes injected with water (top), VR3 A+B- (second from top), VR3 A-B- (3rd from top), and VR3A+B+ (bottom). Oocytes were constantly perfused with Ca²⁺ ND-96 and the solution was heated by an inline heater device (TC-324B in conjunction with the SH-27A in line heater; Warner Instrument Corp.). Ramp induced currents (in microAmperes, as indicated on the y-axis) obtained at temperatures from 37 deg C to 46 deg C are displayed; only current traces at the higher temperatures are labeled with the corresponding temperature.

Perfusion induced currents ($I_{\text{perfusion}}$; Figure 11).

The onset of bath perfusion elicited an increase in membrane conductance in oocytes that had been injected with RNA transcribed from the cloned human VR1 with and without RNA transcribed from VR3A+B- receptor cDNA as shown in FIGURE 11. An oocyte injected with VR1 cRNA (a) was challenged with voltage ramps between -120 and $+80$ mV over 400 msec. The ramp-induced currents were increased after onset of perfusion of the oocyte at a rate of 10 ml/min. Control ramp induced currents from oocytes in still extracellular saline (C) were recorded prior to the onset of bath perfusion of Ca^{2+} ND-96. During perfusion, the ramp induced currents increased, indicating an increase in conductance (P). Subsequent perfusion of 10 μM ruthenium red (RR) did not block the perfusion induced current (P) in VR1-expressing oocytes. Thus, the perfusion-induced current observed in oocytes expressing VR1 alone was insensitive to 10 μM ruthenium red (A, "RR"). However, the perfusion-induced current elicited in oocytes expressing VR3A+B- and VR1 were dramatically inhibited by 10 μM ruthenium red (B, "RR"). This difference was observed in 6 oocytes. The block was partially reversible (not shown). (b). An oocyte injected with VR1 cRNA together with VR3 A+B- was challenged with voltage ramps between -120 and $+80$ mV over 400 msec. The ramp-induced currents were increased after onset of perfusion of the oocyte at a rate of 10 ml/min. Perfusion activated currents had similar magnitudes in both sets of oocytes. V_{rev} for the RR inhibited current was about -13 mV (arrow). The currents induced by both agonists were strongly outwardly rectifying, as reported previously for the rat VR1 (Caterina et al., 1997).

Capsaicin induced currents in oocytes expressing human VR1 in the presence or absence of human VR3 A+B- (Figure 12).

Currents elicited by 1 μ M capsaicin (FIGURE 12) were measured at a holding voltage of +50 mV since responses were largest at this voltage due to profound outward rectification of the capsaicin-induced currents (Caterina et al., 1997). Shown are 3 examples of oocytes injected with VR1 and water (a control for VR3 injection) (a) and oocytes injected with VR1 cRNA and VR3 A+B- cRNA (b). Capsaicin was bath applied during the time indicated by the solid horizontal bar. The magnitude of the primary response to 1 μ M capsaicin was diminished when VR1 cRNA was co-expressed with VR3 A+B- cRNA at equal ratios (2.9 ng each). The most notable difference was the magnitude and decay rate of the secondary response, the second hump that was usually obtained at the beginning of the washout of capsaicin.

EXAMPLE 6 Characterization of Human VR3 in mammalian cell lines.

Human HEK293, CHO-K1 and COS-7 cells are transfected with human VR3 isoforms pVR3A+B-R, pVR3A-B-R, or pVR3A+B+R,. Transient transfections 1 μ g of pVR3R per 10^6 cells per 100 mm dish are performed using the Effectene tranfection kit (Quiagen; 301425). Three days after transfection, cells are plated onto 96-well plates (Biocoat, poly-D-lysine coated black/clear plate; Becton Dickinson part # 354640). After one day, wells are rinsed with F12/DMEM, then incubated in Fluo-4 (2 μ M) with Pluronic F-127 (20%, 40 μ l used in 20 mls total volume) for 1 hour at room temperature. Plates are assayed using the FLIPR (Molecular Devices, FL-101). Cells are challenged with solutions of different osmolarity (40 μ l added to 80 μ l at a velocity of 50 μ l/sec). Some wells are vigorously mixed to simulate increased perfusion of the cells with extracellular solution. Transfections with vector alone are used as controls.

After three days the cells are selected in the presence of neomycin (200 μ g/ml) and grown through three 1:10 dilution passages for approximately two weeks. Individual colonies are picked and grown in 6-well dishes. Cells are then plated onto 96-well plates (Biocoat, poly-D-lysine coated black/clear plate; Becton Dickinson part # 354640) and

grown to confluence for three days. Wells are rinsed with F12/DMEM, then incubated in Fluo-4 (2 μ M) with Pluronic acid (20%, 40 μ l used in 20 mls total volume) for 1 hour at room temperature. Plates are assayed using the FLIPR (Molecular Devices, FL-101). Cells are challenged with agonists (at 3-fold concentration in 40 μ l added to 80 μ l at a velocity of 50 μ l/sec).

The whole cell patch clamp technique (Hamill et al., 1981) is used to record ligand-induced currents from HEK293 stably expressing human VR1 receptor maintained for >2 days on 12 mm coverslips. Cells are visualized using a Nikon Diaphot 300 with DIC Nomarski optics. Cells are continuously perfused in a physiological saline (~0.5 ml/min) unless otherwise indicated. The standard physiological saline ("Tyrodes") contains: 130 mM NaCl, 4 mM KCl, 1 mM CaCl₂, 1.2mM MgCl₂, and 10mM hemi-Na-HEPES (pH 7.3, 295-300 mOsm as measured using a Wescor 5500 vapor-pressure (Wescor, Inc., Logan, UT). Recording electrodes are fabricated from borosilicate capillary tubing (R6; Garner Glass, Claremont, CA), the tips are coated with dental periphery wax (Miles Laboratories, South Bend, IN), and have resistances of 1-2 M Ω when containing intracellular saline: 100 mM K-gluconate, 25 mM KCl, 0.483 mM CaCl₂, 3 mM MgCl₂, 10 mM hemi-Na-HEPES and 1 mM K₄-BAPTA (100nM free Ca⁺²); pH 7.4, with dextrose added to achieve 290 mOsm). Liquid junction potentials are -18 mV using standard pipette and bath solutions as determined both empirically and using the computer program JPCalc ((Barry, 1994)). All voltages shown are corrected for liquid junction potential. Current and voltage signals are detected and filtered at 2 kHz with an Axopatch 1D patch-clamp amplifier (Axon Instruments, Foster City, CA), digitally recorded with a DigiData 1200B laboratory interface (Axon Instruments), and PC compatible computer system and stored on magnetic disk for off-line analysis. Data acquisition and analysis are performed with PClamp software. Slow changes in holding current are detected and filtered at 2 kHz, and recorded with a LPF202A DC amplifier (Warner, Hamden, CT) and VR10B digital data recorder (Instrutech, Great Neck, NY) onto video tape. The signal is later analyzed at 10 Hz using Axotape software.

The total membrane capacitance (C_m) is determined as the difference between the maximum current after a 30 mV hyperpolarizing voltage ramp from -68 mV generated at a rate of 10 mV/ms and the steady state current at the final potential (-98 mV) (Dubin et al., 1999).

Apparent reversal potentials (V_{rev}) of ligand-induced conductance changes are determined using a voltage-ramp protocol (Dubin et al., 1999). Voltage ramps are applied every 1 second and the resulting whole cell ramp-induced currents were recorded. Usually the voltage was ramped from negative to positive to negative values. The current required to clamp the cells at -68 mV is continuously monitored. Ligand-induced conductances are determined from whole-cell currents elicited by a voltage-ramp protocol in the presence and absence of ligand. Voltage ramp-induced currents measured before (control) and in the presence of ligand are compared to reveal the effect of the ligand on the channel to modulate the channel current output. The voltage at which there is no net ligand-induced current is determined (V_{rev}).

EXAMPLE 7 -Primary Structure Of The Human VR3 receptor Protein

The present invention describes 3 isoforms of hVR3: A+B-, A-B-, and A+B+. The nucleotide sequences of human VR3 receptor cDNAs revealed single large open reading frame of about 2616, 2436 and 2229 base pairs encoding 871, 811, and 742 amino acids for human VR3 A+B-, A-B- and A+B+, respectively. The cDNA for VR3 A+B- has 5' and 3'-untranslated extensions of about 337 and about 547 nucleotides. The cDNA for VR3 A+B+ has 5' and 3'-untranslated extensions of about 836 and about 994 nucleotides. The first in-frame methionine was designated as the initiation codon for an open reading frame that predicts human VR3 receptor proteins with an estimated molecular mass (M_r) of about 98,242 Da, 91,294 Da and 83,310 Da for the isoforms A+B-, A-B- and A+B+, respectively. The A+B- isoform encodes a protein of 871 amino acids. The VR3 A-B- contains a deletion of 60 amino acids from amino acid 382 to 441 of VR3A+B-. VR3 A+B+ is identical to A+B-

until amino acid 736 after which there are 6 divergent amino acids and a stop codon. The VR3 A+B+ isoform extends 20 amino acids after the putative TM6.

The predicted human VR3 receptor proteins were aligned with nucleotide and protein databases and found to be related to the vanilloid receptor family (VR1 and VR2). There are several conserved motifs found in this family of receptor including a large putative N-terminal hydrophilic segment (about 467 amino acids), three putative ankyrin repeat domains in the N-terminus region, 6 predicted transmembrane regions and a pore region. VR3 A+B- is 43 % identical to human VR1, 39% identical to both human VRL-1 (AF129112) and human VRL (AF103906). Thus the VR3 receptor described herein is clearly a novel gene of the vanilloid receptor family.

VR3A+B- and VR3A-B- forms are very similar to the rat stretch-inhibitable channel SIC [genbank accession AB015231] from amino acid 694 to the end. SIC, encoded by 529 amino acids, is thought to form an ion channel inhibited by stretch. It lacks the large N-terminal cytoplasmic domain of the VR family but contains a sequence homologous to the A exon prior to the putative TM1.

The complete genomic sequence of the VR3 coding regions described herein appears to be found in a 380512 base pair sequence submission to genbank (homo sapiens clone RPCII-7G5 (AC007834), direct submission by Worley,K.C.). This genbank entry list many fragments of DNA sequence and a proposed contiguous sequence, but lacks any analysis of the nucleic acid sequence and fails to characterize the features of the VR3 nucleic acid sequences, or describe the presence of the VR3 gene.

- 1 Comparison of the sequences the present invention and the genomic sequence reveal that VR3 gene is composed of at least 15 exons. "A" is a 60 amino acid insert in the putative N-terminal cytoplasmic domain that found in cDNAs obtained from both pituitary and prostate cDNA libraries. There are intron/exon border sequences at the A and B inserts.

The A+B- and A+B+ isoforms contain a domain in the N-terminal putative cytoplasmic region with homology to ankyrin repeat domain consensus sequences.

Thus, the VR3 A+ isoforms appear to have a similar architecture as that predicted for VR1. The first domain was significantly similar to the consensus sequence ($E = 2.6 \times 10^{-5}$). The next 2 domains are not significant taken by themselves ($e = 37$ and $E = 2.7$) however taken as a whole, this region is likely to contain 3 ankyrin binding domains. The A- isoform is missing 20 amino acids of the putative 3rd ankyrin repeat domain and the juxtaposed sequence does not conform to an ankyrin repeat domain.

The VR3 A+B- isoform has two potential myristoylation sites: [GPGGE] in N terminal and between TM4 and TM5. Putative phosphorylation sites include: 7 potential PKC phosphorylation sites: T112, S134, T175, T190, T380, S403, S688 [however, S688 is in a putative extracellular region]; 1 potential PKA and PKG phosphorylation site: T181; 12 potential casein kinase II phosphorylation sites: T89, S162, T181, T395, S416, S422, S432, T426, S432, S441, T740, S836; 3 potential mammary gland casein kinase phosphorylation sites (SxE): S4, S726, S758; 1 potential tyrosine kinase phosphorylation site: Y411 [present in the "A" insert]; and 1 potential N-linked glycosylation site: N651, [N201, N207 are in the putative intracellular N-terminal domain and are unlikely to be glycosylated]. There are no putative CaM binding domains in any isoform described in the present invention.

In the A- isoform lacking the "A" insert 2 PKC phosphorylation sites, 6 casein kinase II phosphorylation sites, and the only putative tyrosine kinase phosphorylation site is lacking.

The A+B+ isoform encodes a putative protein that lacks 2 casein kinase II phosphorylation sites in the C-terminal putative intracellular region.

EXAMPLE 8- Expression of human VR3 in tissues.

Expression using a DNA array (Luo et al., 1999). DNA array distribution analysis indicates that human VR3 mRNAs were expressed in a variety of tissues, and there was some overlap of expression with VR1 at a whole tissue level [Figure 13]. The cRNA

from the tissue type indicated in the left column contained sequences encoding the human VR3 (middle column). Expression of human VR1 is shown in the right column and in most cases overlapped with VR3 expression. NOTE: NS was non-significant expression of human VR1. The sequence of human VR3 DNA immobilized on the DNA array was (SEQ.ID.NO.:17)

```
CCACCATCCTGGACATTGAGCGCTCCTTCCCCGTATTCCTGAGGAAGGCCTTC
CGCTCTGGGGAGATGGTCACCGTGGGCAAGAGCTCGGACGGCACTCCTGACC
GCAGTGGTGCTTCAGGGTGGATGAGGTGAACTGGTCTCACTGGAACCAGAAC
TTGGGCATCATCAACGAGGACCCGGGCAAGAATGAGACCTACCAGTATTATG
G CTTCTCGCATACCGTGGGCGGCC.
```

This DNA sequence is not present in A+B+, only A+B- and A-B- isoforms. Note that the cDNA species was cloned from prostate gland (*; Figure 13).

Northern blot analysis revealed expression of VR3 in whole brain, placenta, lung, kidney, pancreas and prostate.

EXAMPLE 9-Cloning human VR3 receptor cDNA into E. coli Expression Vectors

Recombinant human VR3 receptor is produced in E. coli following the transfer of the human VR3 receptor expression cassette into E. coli expression vectors, including but not limited to, the pET series (Novagen). The pET vectors place human VR3 receptor expression under control of the tightly regulated bacteriophage T7 promoter. Following transfer of this construct into an E. coli host that contain a chromosomal copy of the T7 RNA polymerase gene driven by the inducible lac promoter, expression of human VR3 receptor is induced when an appropriate lac substrate (IPTG) is added to the culture. The levels of expressed human VR3 receptor are determined by the assays described herein.

The cDNA encoding the entire open reading frame for human VR3 receptor is inserted into the NdeI site of pET [16]11a. Constructs in the positive orientation are

identified by sequence analysis and used to transform the expression host strain BL21. Transformants are then used to inoculate cultures for the production of human VR3 receptor protein. Cultures may be grown in M9 or ZB media, whose formulation is known to those skilled in the art. After growth to an $OD_{600} = 1.5$, expression of human VR3 receptor is induced with 1 mM IPTG for 3 hours at 37°C.

EXAMPLE 10-Cloning human VR3 receptor cDNA into a Baculovirus Expression Vector for Expression in Insect Cells

Baculovirus vectors, which are derived from the genome of the AcNPV virus, are designed to provide high level expression of cDNA in the Sf9 line of insect cells (ATCC CRL# 1711). Recombinant baculoviruses expressing human VR3 receptor cDNA is produced by the following standard methods (InVitrogen Maxbac Manual): the human VR3 receptor cDNA constructs are ligated into the polyhedrin gene in a variety of baculovirus transfer vectors, including the pAC360 and the BlueBac vector (InVitrogen). Recombinant baculoviruses are generated by homologous recombination following co-transfection of the baculovirus transfer vector and linearized AcNPV genomic DNA [Kitts, P.A., Nuc. Acid. Res. 18: 5667 (1990)] into Sf9 cells. Recombinant pAC360 viruses are identified by the absence of inclusion bodies in infected cells and recombinant pBlueBac viruses are identified on the basis of β -galactosidase expression (Summers, M. D. and Smith, G. E., Texas Agriculture Exp. Station Bulletin No. 1555). Following plaque purification, human VR3 receptor expression is measured by the assays described herein.

The cDNA encoding the entire open reading frame for human VR3 receptor is inserted into the BamHI site of pBlueBacII. Constructs in the positive orientation are identified by sequence analysis and used to transfect Sf9 cells in the presence of linear AcNPV mild type DNA.

Authentic, active human VR3 receptor is found in the cytoplasm of infected cells. Active human VR3 receptor is extracted from infected cells by hypotonic or detergent lysis.

EXAMPLE 11-Cloning human VR3 receptor cDNA into a yeast expression vector

Recombinant human VR3 receptor is produced in the yeast *S. cerevisiae* following insertion of the optimal human VR3 receptor cDNA cistron into expression vectors designed to direct the intracellular or extracellular expression of heterologous proteins. In the case of intracellular expression, vectors such as EmBLyex4 or the like are ligated to the human VR3 receptor cistron [Rinas, U. *et al.*, Biotechnology 8: 543-545 (1990); Horowitz B. *et al.*, J. Biol. Chem. 265: 4189-4192 (1989)]. For extracellular expression, the human VR3 receptor cistron is ligated into yeast expression vectors which fuse a secretion signal (a yeast or mammalian peptide) to the NH₂ terminus of the human VR3 receptor protein [Jacobson, M. A., Gene 85: 511-516 (1989); Riett L. and Bellon N. Biochem. 28: 2941-2949 (1989)].

These vectors include, but are not limited to pAVE1>6, which fuses the human serum albumin signal to the expressed cDNA [Steep O. Biotechnology 8: 42-46 (1990)], and the vector pL8PL which fuses the human lysozyme signal to the expressed cDNA [Yamamoto, Y., Biochem. 28: 2728-2732)]. In addition, human VR3 receptor is expressed in yeast as a fusion protein conjugated to ubiquitin utilizing the vector pVEP [Ecker, D. J., J. Biol. Chem. 264: 7715-7719 (1989), Sabin, E. A., Biotechnology 7: 705-709 (1989), McDonnell D. P., Mol. Cell Biol. 9: 5517-5523 (1989)]. The levels of expressed human VR3 receptor are determined by the assays described herein.

EXAMPLE 12-Purification of Recombinant human VR3 receptor

Recombinantly produced human VR3 receptor may be purified by antibody affinity chromatography.

Human VR3 receptor antibody affinity columns are made by adding the anti-human VR3 receptor antibodies to Affigel-10 (Bio-Rad), a gel support that is pre-activated with N-hydroxysuccinimide esters such that the antibodies form covalent linkages with the agarose gel bead support. The antibodies are then coupled to the gel via amide bonds with the spacer arm. The remaining activated esters are then quenched with 1M ethanolamine HCl (pH 8). The column is washed with water followed by 0.23 M glycine HCl (pH 2.6) to remove any non-conjugated antibody or extraneous protein. The column is then equilibrated in phosphate buffered saline (pH 7.3) together with appropriate membrane solubilizing agents such as detergents and the cell culture supernatant or cell extract containing solubilized human VR3 receptor is slowly passed through the column. The column is then washed with phosphate-buffered saline together with detergents until the optical density (A280) falls to background, then the protein is eluted with 0.23 M glycine-HCl (pH 2.6) together with detergents. The purified human VR3 receptor protein is then dialyzed against phosphate buffered saline.

References

Bevan, S., Hothi, S., Hughes, G., James, I. F., Rang, H. P., Shah, K., Walpole, C. S. J., and Yeats, J. C. (1992). Capsazepine: a competitive antagonist of the sensory neuron excitant capsaicin. *Br. J. Pharmacol.* 107, 544-52.

Bevan, S., and Szolcsanyi, J. (1990). Sensory neuron-specific actions of capsaicin: mechanisms and applications. *Trends Pharmacol. Sci.* 11, 330-3.

Blackstone, C., and Sheng, M. (1999). Protein targeting and calcium signaling microdomains in neuronal cells. *Cell Calcium* 26, 181-192.

Caterina, M. J., Rosen, T. A., Tominaga, M., Brake, A. J., and Julius, D. (1999). A capsaicin-receptor homolog with a high threshold for noxious heat. *Nature* (London) 398, 436-441.

Caterina, M. J., Schumacher, M. A., Tominaga, M., Rosen, T. A., Levine, J. D., and Julius, D. (1997). The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* (London) 389, 816-824.

Fraser, S. P., Moon, C., and Djamgoz, M. B. A. (1993). Electrophysiology of *Xenopus oocytes*: An expression system in molecular neurobiology., D. Wallis, ed. (Oxford, UK: IRL).

Gupta, P. D., and Pushkala, K. (1999). Importance of the role of calcium in programmed cell death: a review. *Cytobios* 99, 83-95.

Lu, M., and Wong, F. (1987). A DNA deletion associated with multiple impaired transcripts in the visual mutant *trp*. *Invest. Ophthalmol. Visual Sci.* 28, 2092-5.

Luo, L., Salunga, R. C., Guo, H., Bitner, A., Joy, K. C., Galindo, J. E., Xiao, H., Rogers, K. E., Wan, J. S., Jackson, M. R., and Erlander, M. G. (1999). Gene expression profiles of laser-captured adjacent neuronal subtypes. *Nat. Med.* (N. Y.) 5, 117-121.

Minke, B., and Selinger, Z. (1996). The roles of trp and calcium in regulating photoreceptor function in *Drosophila*. In *Curr. Opin. Neurobiol.*, pp. 459-466.

Oh, U., Hwang, S. W., and Kim, D. (1996). Capsaicin activates a nonselective cation channel in cultured neonatal rat dorsal root ganglion neurons. *J. Neurosci.* 16, 1659-67.

Suzuki, M., Sato, J., Kutsuwada, K., Ooki, G., and Imai, M. (1999). Cloning of a stretch-inhibitable nonselective cation channel. *J. Biol. Chem.* 274, 6330-6335.

Szallasi, A. (1995). Autoradiographic visualization and pharmacological characterization of vanilloid (capsaicin) receptors in several species, including man. *Acta Physiol. Scand., Suppl.* 155, 1-68.

Szolcsanyi, J. (1993). Actions of capsaicin on sensory receptors. In Capsaicin Study Pain, J. N. Wood, ed.: Academic, London, UK), pp. 1-26.

Szolcsanyi, J. (1996). Capsaicin-sensitive sensory nerve terminals with local and systemic efferent functions: facts and scopes of an unorthodox neuroregulatory mechanism. *Prog. Brain Res.* 113, 343-359.

Szolcsanyi, J., Szallasi, A., Szallasi, Z., Joo, F., and Blumberg, P. M. (1991). Resiniferatoxin. An ultrapotent neurotoxin of capsaicin-sensitive primary afferent neurons. *Ann. N. Y. Acad. Sci.* 632, 473-5.

Tominaga, M., Caterina, M. J., Malmberg, A. B., Rosen, T. A., Gilbert, H., Skinner, K., Raumann, B., Basbaum, A. I., and Julius, D. (1998). The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron* 21, 531-543.

van Haasteren, G., Li, S., Muda, M., Susini, S., and Schlegel, W. (1999). Calcium signalling and gene expression. In *J. Recept. Signal Transduction Res.*, pp. 481-492.

Wall, P. D., and Melzack, R. (1994). Textbook of Pain (New York: Churchill Livingstone).

Wood, J. N., Winter, J., James, I. F., Rang, H. P., Yeats, J., and Bevan, S. (1988). Capsaicin-induced ion fluxes in dorsal root ganglion cells in culture. *J. Neurosci.* 8, 3208-20.

WHAT IS CLAIMED IS:

1. An isolated and purified DNA molecule which encodes a human VR3 receptor protein.
2. The isolated and purified DNA molecule of claim 1, having a nucleotide sequence selected from a group consisting of: (SEQ.ID.NO.:5); (SEQ.ID.NO.:6); (SEQ.ID.NO.:8); (SEQ.ID.NO.:10) and (SEQ.ID.NO.:11)
3. The isolated and purified DNA molecule of claim 1, wherein said DNA molecule is genomic DNA.
4. An expression vector for expression of an human VR3 receptor protein in a recombinant host, wherein said vector comprises a DNA sequence encoding human VR3 receptor protein.
5. The expression vector of claim 4, wherein the DNA sequence encoding human VR3 receptor protein is selected from a group consisting of: (SEQ.ID.NO.:5); (SEQ.ID.NO.:6); (SEQ.ID.NO.:8); (SEQ.ID.NO.:10) and (SEQ.ID.NO.:11).
6. The expression vector of claim 4, wherein the expression vector contains genomic DNA encoding human VR3 receptor protein.
7. A recombinant host cell comprising the expression vector of claim 4.

8. The recombinant host cell of claim 7, wherein said expression vector comprises a nucleotide sequence selected from a group consisting of: (SEQ.ID.NO.:5); (SEQ.ID.NO.:6); (SEQ.ID.NO.:8); (SEQ.ID.NO.:10) and (SEQ.ID.NO.:11).

9. The recombinant host cell of claim 7, wherein said expression vector comprises genomic DNA encoding a human VR3 receptor.

10. A protein, in substantially pure form which functions as human VR3 receptor protein.

11. The protein according to claim 10, having an amino acid sequence selected from a group consisting of: (SEQ.ID.NO.:7) (SEQ.ID.NO.:9) and (SEQ.ID.NO.:12).

12. A monospecific antibody immunologically reactive with human VR3 receptor protein.

13. The antibody of Claim 12, wherein the antibody blocks activity of the human VR3 receptor.

14. A process for expression of human VR3 receptor protein in a recombinant host cell comprising:

- (a) transferring the expression vector of Claim 4 into suitable host cells; and
- (b) culturing the host cells of step (a) under conditions which allow expression of the human VR3 receptor protein from the expression vector.

15. A method of identifying compounds that modulate human VR3 receptor protein activity, comprising:

- (a) combining a modulator of human VR3 receptor protein activity with human VR3 receptor protein. and
- (b) measuring an effect of the modulator on the human VR3 receptor protein.

16. The method of claim 15, wherein the effect of the modulator on the protein is inhibiting or enhancing binding of human VR3 receptor ligands.

17. The method of claim 15, wherein the effect of the modulator on the human VR3 receptor protein is stimulation or inhibition of human VR3 receptor activity protein.

18. A compound active in the method of Claim 15.

19. A compound active in the method of Claim 15, wherein said compound is an agonist or antagonist of a human VR3 receptor.

20. A compound active in the method of Claim 15, wherein said compound is a modulator of expression of a human VR3 receptor.

21. A pharmaceutical composition comprising a compound active in the method of Claim 15.

22. A method of treating a patient in need of such treatment for a condition which is mediated by a human VR3 receptor, comprising administration of a compound active in the method of Claim 15.

1/18

FIGURE 1

SEQ.ID.NO.5. Human VR3A+B- nucleotide sequence of the coding sequence (2616 bp).

ATGGCGGATTCCAGCGAAGGCCCGCGCGGGGGCCCGGGGAGGTGGCTGAG
CTCCCCGGGGATGAGAGTGGCACCCAGGTGGGGAGGCTTTTCCTCTCTCCTC
CCTGGCCAATCTGTTTGAGGGGGAGGATGGCTCCCTTTTCGCCCTCACCGGCTG
ATGCCAGTCGCCCTGCTGGCCCAGGCGATGGGCGACCAAATCTGCGCATGAA
GTTCCAGGGCGCCTTCCGCAAGGGGGTGCCCAACCCCATCGATCTGCTGGAG
TCCACCCTATATGAGTCCTCGGTGGTGCCTGGGCCCCAAGAAAGCACCCATGG
ACTCACTGTTTGACTACGGCACCTATCGTCACCACTCCAGTGACAACAAGAG
GTGGAGGAAGAAGATCATAGAGAAGCAGCCGCAGAGCCCCAAAGCCCCTGC
CCCTCAGCCGCCCCCATCCTCAAAGTCTTCAACCGGCCTATCCTCTTTGACA
TCGTGTCCCGGGGCTCCACTGCTGACCTGGACGGGCTGCTCCCATTTCTTGCTG
ACCCACAAGAAACGCCTAACTGATGAGGAGTTTCGAGAGCCATCTACGGGGA
AGACCTGCCTGCCCAAGGCCTTGCTGAACCTGAGCAATGGCCGCAACGACAC
CATCCCTGTGCTGCTGGACATCGCGGAGCGCACCGGCAACATGCGGGAGTTC
ATTAACCTCGCCCTTCCGTGACATCTACTATCGAGGTCAGACAGCCCTGCACAT
CGCCATTGAGCGTCGCTGCAAACACTACGTGGAACCTTCTCGTGGCCCAGGGA
GCTGATGTCCACGCCAGGCCCGTGGGCGCTTCTTCCAGCCCAAGGATGAGG
GGGGCTACTTCTACTTTGGGGAGCTGCCCTGTGCTGGCTGCCTGCACCAAC
CAGCCCCACATTGTCAACTACCTGACGGAGAACCCCCACAAGAAGGCGGACA
TGCGGCGCCAGGACTCGCGAGGCAACACAGTGCTGCATGCGCTGGTGGCCAT
TGCTGACAACACCCGTGAGAACACCAAGTTTGTTACCAAGATGTACGACCTG
CTGCTGCTCAAGTGTGCCCGCCTCTTCCCCGACAGCAACCTGGAGGCCGTGCT
CAACAACGACGGCCTCTCGCCCCTCATGATGGCTGCCAAGACGGGCAAGATT
GGGATCTTTCAGCACATCATCCGGCGGGAGGTGACGGATGAGGACACACGGC
ACCTGTCCCGCAAGTTCAAGGACTGGGCCTATGGGCCAGTGTATTCCTCGCTT
TATGACCTCTCCTCCCTGGACACGTGTGGGGAAGAGGCCTCCGTGCTGGAGA
TCCTGGTGTACAACAGCAAGATTGAGAACCGCCACGAGATGCTGGCTGTGGA
GCCCATCAATGAACTGCTGCGGGACAAGTGGCGCAAGTTCGGGGCCGTCTCC
TTCTACATCAACGTGGTCTCCTACCTGTGTGCCATGGTCATCTTCACTCTCACC
GCCTACTACCAGCCGCTGGAGGGGCACACCGCCGTACCCTTACCGCACACGG
TGGACTACCTGCGGCTGGCTGGCGAGGTCATTACGCTCTTCACTGGGGTCCTG
TTCTTCATCACCACATCAAAGACTTGTTTCATGAAGAAATGCCCTGGAGTGA
ATTCTCTCTTCATTGATGGCTCCTTCCAGCTGCTCTACTTCATCTACTCTGTCC
TGGTGATCGTCTCAGCAGCCCTCTACCTGGCAGGGATCGAGGCCTACCTGGC
CGTGATGGTCTTTGCCCTGGTCCTGGGCTGGATGAATGCCCTTTACTTCACCC
GTGGGCTGAAGCTGACGGGGACCTATAGCATCATGATCCAGAAGATTCTCTT

2/18

CAAGGACCTTTTCCGATTCCTGCTCGTCTACTTGCTCTTCATGATCGGCTACGC
TTCAGCCCTGGTCTCCCTCCTGAACCCGTGTGCCAACATGAAGGTGTGCAATG
AGGACCAGACCAACTGCACAGTGCCCACTTACCCCTCGTGCCGTGACAGCGA
GACCTTCAGCACCTTCCTCCTGGACCTGTTTAAGCTGACCATCGGCATGGGCG
ACCTGGAGATGCTGAGCAGCACCAAGTACCCCGTGGTCTTCATCATCCTGCT
GGTGACCTACATCATCCTCACCTTTGTGCTGCTCCTCAACATGCTCATTGCCC
TCATGGGCGAGACAGTGGGCCAGGTCTCCAAGGAGAGCAAGCACATCTGGA
AGCTGCAGTGGGCCACCACCATCCTGGACATTGAGCGCTCCTTCCCCGTATTC
CTGAGGAAGGCCTTCCGCTCTGGGGAGATGGTCACCGTGGGCAAGAGCTCGG
ACGGCACTCCTGACCGCAGGTGGTGCTTCAGGGTGGATGAGGTGAACTGGTC
TCACTGGAACCAGAACTTGGGCATCATCAACGAGGACCCGGGCAAGAATGA
GACCTACCAGTATTATGGCTTCTCGCATACCGTGGGCCGCCTCCGCAGGGATC
GCTGGTCCTCGGTGGTACCCCGCGTGGTGGAAGTGAACAAGAACTCGAACCC
GGACGAGGTGGTGGTGCCTCTGGACAGCATGGGGAACCCCCGCTGCGATGGC
CACCAGCAGGGTTACCCCCGCAAGTGGAGGACTGATGACGCCCCGCTCTAG

3/18

FIGURE 2

SEQ.ID.NO.6. The nucleotide sequence of human VR3A+B- is shown including 337 bp 5' UT and 547 bp 3'UT .

CAATTGGGATTTAAACCCAGGGACTATCCAGCCCCAAAGCCCTTCCCACCAC
ACCAGGTGGCCTGTCCTGGGGCCAGCTCTGCACACAGGGCCTGGTGCCCCCG
GGGTGCTTGGGAAGTGGCAGGGCAGAGGTGGGCCCTGTGGCTGTTCTGGCTC
AGCTTCTAAAACAAGAGCCTCTGCTGGGGGCAGAGGGGCCGTGAACCCCTGA
AATGTTAGGCAGATACCCTGTGGGAGCTTTGTTCTGGGATGCTAAGAACCGC
TTGAGGATTTAAGCTTTGCCACTTTGGCTCCGGAGCAAGGGCAGAGGCTGAG
CAGTGCAGACGGGCCTGGGGCAGGCATGGCGGATTCCAGCGAAGGCCCCCG
CGCGGGGCCCCGGGAGGTGGCTGAGCTCCCCGGGGATGAGAGTGGCACCCC
AGGTGGGGAGGCTTTTCTCTCTCCTCCCTGGCCAATCTGTTTGAGGGGGAGG
ATGGCTCCCTTTCGCCCTCACCGGCTGATGCCAGTCGCCCTGCTGGCCCAGGC
GATGGGCGACCAAATCTGCGCATGAAGTTCCAGGGCGCCTTCCGCAAGGGGG
TGCCCAACCCCATCGATCTGCTGGAGTCCACCCTATATGAGTCCTCGGTGGTG
CCTGGGCCCCAAGAAAGCACCCATGGACTCACTGTTTGACTACGGCACCTATC
GTCACCACTCCAGTGACAACAAGAGGTGGAGGAAGAAGATCATAGAGAAGC
AGCCGCAGAGCCCCAAAGCCCCTGCCCTCAGCCGCCCCCATCCTCAAAGT
CTTCAACCGGCCTATCCTCTTTGACATCGTGTCCCGGGGCTCCACTGCTGACC
TGGACGGGCTGCTCCCATTTCTTGCTGACCCACAAGAAACGCCTAACTGATGA
GGAGTTTTCGAGAGCCATCTACGGGGAAGACCTGCCTGCCCAAGGCCTTGCTG
AACCTGAGCAATGGCCGCAACGACACCATCCCTGTGCTGCTGGACATCGCGG
AGCGCACCGGCAACATGCGGGAGTTCATTAACCTCGCCCTTCCGTGACATCTA
CTATCGAGGTCAGACAGCCCTGCACATCGCCATTGAGCGTCGCTGCAAACAC
TACGTGGAACCTTCTCGTGGCCCAGGGAGCTGATGTCCACGCCCAGGCCCGTG
GGCGCTTCTTCCAGCCCAAGGATGAGGGGGGGCTACTTCTACTTTGGGGAGCT
GCCCCTGTCGCTGGCTGCCTGCACCAACCAGCCCCACATTGTCAACTACCTGA
CGGAGAACCCCCACAAGAAGGCGGACATGCGGCGCCAGGACTCGCGAGGCA
ACACAGTGCTGCATGCGCTGGTGGCCATTGCTGACAACACCCGTGAGAACAC
CAAGTTTGTACCAAGATGTACGACCTGCTGCTGCTCAAGTGTGCCCGCCTCT
TCCCCGACAGCAACCTGGAGGCCGTGCTCAACAACGACGGCCTCTCGCCCCCT
CATGATGGCTGCCAAGACGGGCAAGATTGGGATCTTTCAGCACATCATCCGG
CGGGAGGTGACGGATGAGGACACACGGCACCTGTCCCGCAAGTTCAAGGAC
TGGGCCTATGGGCCAGTGTATTCCTCGCTTTATGACCTCTCCTCCCTGGACAC
GTGTGGGGAAAGAGGCCTCCGTGCTGGAGATCCTGGTGTACAACAGCAAGATT
GAGAACCGCCACGAGATGCTGGCTGTGGAGCCCATCAATGAAGTGTGCGGG
ACAAGTGGCGCAAGTTCGGGGCCGTCTCCTTCTACATCAACGTGGTCTCCTAC
CTGTGTGCCATGGTCATCTTCACTCTCACCGCCTACTACCAGCCGCTGGAGGG
CACACCGCCGTACCCTTACCGCACACGGTGGACTACCTGCGGCTGGCTGGC

4/18

GAGGTCATTACGCTCTTCACTGGGGTCCTGTTCTTCATCACCAACATCAAAGA
CTTGTTTCATGAAGAAATGCCCTGGAGTGAATTCTCTCTTCATTGATGGCTCCT
TCCAGCTGCTCTACTTCATCTACTCTGTCTGGTGATCGTCTCAGCAGCCCTCT
ACCTGGCAGGGATCGAGGCCTACCTGGCCGTGATGGTCTTTGCCCTGGTCCTG
GGCTGGATGAATGCCCTTTACTTCACCCGTGGGCTGAAGCTGACGGGGACCT
ATAGCATCATGATCCAGAAGATTCTCTTCAAGGACCTTTTCCGATTCTGCTC
GTCTACTTGCTCTTCATGATCGGCTACGCTTCAGCCCTGGTCTCCCTCCTGAA
CCCGTGTGCCAACATGAAGGTGTGCAATGAGGACCAGACCAACTGCACAGTG
CCCACTTACCCCTCGTGCCGTGACAGCGAGACCTTCAGCACCTTCCTCCTGGA
CCTGTTTAAGCTGACCATCGGCATGGGCGACCTGGAGATGCTGAGCAGCACC
AAGTACCCCGTGGTCTTCATCATCCTGCTGGTGACCTACATCATCCTCACCTT
TGTGCTGCTCCTCAACATGCTCATTGCCCTCATGGGCGAGACAGTGGGCCAG
GTCTCCAAGGAGAGCAAGCACATCTGGAAGCTGCAGTGGGCCACCACCATCC
TGGACATTGAGCGCTCCTTCCCCGTATTCTGAGGAAGGCCTTCCGCTCTGGG
GAGATGGTCACCGTGGGCAAGAGCTCGGACGGCACTCCTGACCGCAGGTGGT
GCTTCAGGGTGGATGAGGTGAACTGGTCTCACTGGAACCAGAACTTGGGCAT
CATCAACGAGGACCCGGGCAAGAATGAGACCTACCAGTATTATGGCTTCTCG
CATAACCGTGGGCGCCTCCGCAGGGATCGCTGGTCCTCGGTGGTACCCCGCG
TGGTGGAAGTGAACAAGAACTCGAACCCGGACGAGGTGGTGGTGCCTCTGGA
CAGCATGGGGAACCCCCGCTGCGATGGCCACCAGCAGGGTTACCCCCGCAAG
TGGAGGACTGATGACGCCCCGCTCTAGGGACTGCAGCCCAGCCCCAGCTTCT
CTGCCCCACTCATTTCTAGTCCAGCCGCATTTCAAGCAGTGCCTTCTGGGGTGTC
CCCCACACCCTGCTTTGGCCCCAGAGGCGAGGGACCAGTGGAGGTGCCAGG
GAGGCCCCAGGACCCTGTGGTCCCCTGGCTCTGCCTCCCCACCCTGGGGTGG
GGGCTCCCGGCCACCTGTCTTGCTCCTATGGAGTCACATAAGCCAACGCCAG
AGCCCCTCCACCTCAGGCCCCAGCCCCTGCCTCTCCATTATTTATTTGCTCTGC
TCTCAGGAAGCGACGTGACCCCTGCCCCAGCTGGAACCTGGCAGAGGCCTTA
GGACCCCGTTCCAAGTGCACTGCCCGGCCAAGCCCCAGCCTCAGCCTGCGCC
TGAGCTGCATGCGCCACCATTTTTGGCAGCGTGGCAGCTTTGCAAGGGGCTG
GGGCCCTCGGCGTGGGGCCATGCCTTCTGTGTGTTCTGTAGTGTCTGGGATTT
GCCGGTGCTCAATAAATGTTTATTCATTGACGGTGGAAAAAAAAAAAAAA

5/18

FIGURE 3

SEQ.ID.NO.7. Coding sequence for human VR3A+B- (871 amino acids)

MADSSEGPRAGPGEVAELPGDESGTPGGEAFPLSSLANLFEGEDGSLSPSPADAS
RPAGPGDGRPNLRMKFQGAFRKGVPNPIDLLESTLYESSVVPGPCKAPMDSLFD
YGTYRHHSSDNKRWRKKIIEKQPQSPKAPAPQPPILKVFNRPILFDIVSRGSTAD
LDGLLPFLLTTHKKRLTDEEFREPSTGKTCLPKALLNLSNGRNDTIPVLLDIAERTG
NMREFINSPFRDIYYRGQTALHIAIERRCKHYVELLVAQGADVHAQARGRFFQPK
DEGGYFYFGELPLSLAACTNQPHIVNYLTENPHKKADMRRQDSRGNTVLHALV
AIADNTRENTKFVTKMYDLLLLKCARLFPDSNLEAVLNNDGLSPLMMAAKTGKI
GIFQHIIREVTDDEDTRHLSRKFKDWAYGPVYSSLYDLSSLDTCGEEASVLEILVY
NSKIENRHEMLAVEPINELLRDKWRKFGAVSFYINVVSYLCAMVIFTLTAYYQPL
EGTPPYPYRTTVDYLRRLAGEVITLFTGVLFFITNIKDLFMKKCPGVNSLFIDGSFQL
LYFIYSVLVIVSAALYLAGIEAYLAVMVFALVLGWMNALYFTRGLKLTGTYSIMI
QKILFKDLFRFLLVYLLFMIGYASALVSLNPCANMKVCNEDQTNCTVPTYPSCR
DSETFSTFLDLFKLTIGMGDLEMLSSTKYPVVFILLVTYIILTFVLLLNMLIALMGE
TVGQVSKESKHIWKLQWATTILDIERSFPVFLRKAFRSGEMVTVGKSSDGTPDRRWCF
RVDEVNWSHWNQNLGIINEDPGKNETYQYYGFSHTVGRLRRDRWSSVVPVVELNKN
SNPDEVVVPLDSMGNPRCDGHQQGYPRKWRTDDAPL

6/18

FIGURE 4

SEQ.ID.NO.8. Human VR3A-B- nucleotide sequence of the coding sequence (2436 bp).

```
ATGGCGGATTCCAGCGAAGGCCCCGCGCGGGGCCCCGGGAGGTGGCTGAGCTCCCCGGGGATGAGAGTGG
CACCCAGGTGGGGAGGCTTTTCTCTCTCTCTCCCTGGCCAATCTGTTTGAGGGGGAGGATGGCTCCCTTT
CGCCCTCACCGGCTGATGCCAGTCGCCCTGCTGGCCAGGCGATGGGCGACCAAATCTGCGCATGAAGTTC
CAGGGCGCCTTCCGCAAGGGGGTGCCCAACCCCATCGATCTGCTGGAGTCCACCCATATGAGTCCTCGGT
GGTGCCTGGGCCCCAAGAAAGCACCCATGGACTCACTGTTTGACTACGGCACCTATCGTCACCACTCCAGTG
ACAACAAGAGGTGGAGGAAGAAGATCATAGAGAAGCAGCCGAGAGCCCCAAAGCCCCCTGCCCTCAGCCG
CCCCCATCCTCAAAGTCTTCAACCGGCCTATCTCTTTGACATCGTGTCCCGGGGCTCCACTGCTGACCT
GGACGGGCTGCTCCCATTTCTTGCTGACCCACAAGAAACGCCTAACTGATGAGGAGTTTCGAGAGCCATCTA
CGGGGAAGACCTGCCTGCCCAAGGCCTTGCTGAACCTGAGCAATGGCCGCAACGACACCATCCCTGTGCTG
CTGGACATCGCGGAGCGCACCGGCAACATGCGGGAGTTCAATTAACGCCCCCTCCGTGACATCTACTATCG
AGGTGAGACAGCCCTGCACATCGCCATTGAGCGTCGCTGCAAACTACGTGGAACCTTCTCGTGGCCCAGG
GAGCTGATGTCCACGCCAGGCCCGTGGGCGCTTCTTCCAGCCCCAAGGATGAGGGGGGCTACTTCTACTTT
GGGGAGCTGCCCTGTGCTGGCTGCCTGCACCAACAGCCCCACATTGTCAACTACCTGACGGAGAACC
CCACAAGAAGGCGGACATGCGGCGCCAGGACTCGCGAGGCAACACAGTGCTGCATGCGTGGTGGCCATTG
CTGACAACACCCGTGAGAACACCAAGTTTGTACCAAGATGTACGACCTGCTGCTGCTCAAGTGTGCCGCG
CTCTTCCCCGACAGCAACCTGGAGGCCGTGCTCAACAACGACGGCCTCTCGCCCCCTCATGATGGCTGCCAA
GACGGGCAAGATTGAGAACCGCCACGAGATGCTGGCTGTGGAGCCCATCAATGAACCTGCTGCGGGACAAGT
GGCGCAAGTTTCGGGGCCGTCTCCTTCTACATCAACGTGGTCTCCTACCTGTGTGCCATGGTCATCTTCACT
CTCACCGCCTACTACCAGCCGCTGGAGGGCACACCGCCGTACCCTTACCGCACCCACGGTGGACTACCTGCG
GCTGGCTGGCGAGGTCATTACGCTCTTCACTGGGGTCTGTCTTTCATCACCAACATCAAAGACTTGTTC
TGAAGAAATGCCCTGGAGTGAATTCTCTCTTCAATTGATGGCTCCTTCCAGCTGCTCTACTTCTACTCT
GTCCTGGTGATCGTCTCAGCAGCCCTCTACCTGGCAGGGATCGAGGCCTACCTGGCCGTGATGGTCTTTGC
CCTGGTCTTGGGCTGGATGAATGCCCTTTACTTCAACCCGTGGGCTGAAGCTGACGGGGACCTATAGCATCA
TGATCCAGAAGATTCTCTTCAAGGACCTTTTCCGATTCTGCTCGTCTACTTGCTCTTCATGATCGGCTAC
GCTTCAGCCCTGGTCTCCCTCCTGAACCCGTGTGCCAACATGAAGGTGTGCAATGAGGACCAGACCAACTG
CACAGTGCCCACTTACCCCTCGTGCCGTGACAGCGAGACCTTCAGCACCTTCCTCCTGGACCTGTTTAAGC
TGACCATCGGCATGGGCGACCTGGAGATGCTGAGCAGCACCAAGTACCCCGTGGTCTTCATCATCCTGCTG
GTGACCTACATCATCCTCACCTTTGTGCTGCTCCTCAACATGCTCATTGCCCTCATGGGCGAGACAGTGGG
CCAGGTCTCCAAGGAGAGCAAGCACATCTGGAAGCTGCAGTGGGCCACCACCATCCTGGACATTGAGCGCT
CCTTCCCCGTATTCTTGAGGAAGGCCTTCCGCTCTGGGGAGATGGTCAACCGTGGGCAAGAGCTCGGACGGC
ACTCCTGACCGCAGGTGGTGTCTCAGGGTGGATGAGGTGAACTGGTCTCACTGGAACCAGAACTTGGGCAT
CATCAACGAGGACCCGGGCAAGAATGAGACCTACCAGTATTATGGCTTCTCGCATACCGTGGGCGCCCTCC
GCAGGGATCGCTGGTCTCGGTGGTACCCCGCTGGTGGAACTGAACAAGAACTCGAACCCGGACGAGGTG
GTGGTGCCTCTGGACAGCATGGGGAACCCCCGCTGCGATGGCCACCAGCAGGGTTACCCCGCAAGTGGAG
GACTGATGACGCCCCGCTCTAG
```

7/18

FIGURE 5

SEQ.ID.NO.9. Coding sequence for human VR3A-B- (811 amino acids)

MADSSEGPRAGPGEVAELPGDESGTPGGEAFPLSSLANLFEGEDGSLSPSPADASRPAG
PGDGRPNLRMKFQGAFRKGVPNPIDLLESTLYESSVVPGPKKAPMDSLFDYGTyrHHSS
DNKRWRKKIIEKQPQSPKAPAPQPPPIILKVFNRPILFDIVSRGSTADLDGLLPFLLLTHK
KRLTDEEFREPSTGKTCLPKALLNLSNGRNDTIPVLLDIAERTGNMREFINSPFRDIYY
RGQTALHIAIERRCKHYVELLVAQGADVHAQARGRFFQPKDEGGYFYFGELPLSLAACT
NQPHIVNYLTENPHKKADMRRQDSRGNTVLHALVAIADNTRENTKFVTKMYDLLLLKCA
RLFPDSNLEAVLNNDGLSPLMMAAKTGKIENRHEMLAVEPINELLRDKWRKFGAVSFYI
NVVSYLCAMVIFTLTAYYQPLEGTPPYRRTTVDYLRRLAGEVITLFTGVLFITNIKDL
FMKKCPGVNSLFIDGSFQLLYFYISVLVIVSAALYLAGIEAYLAVMVFALVLGWMNALY
FTRGLKLTGTYSIMIQILFKDLFRFLLVYLLFMIGYASALVSLNPNPCANMKVCNEDQT
NCTVPTYPSCRDSETFSTFLLDLFKLTIGMGDLEMLSSTKYPVVFIIILLVTYIILTFVL
LLNMLIALMGETVGQVSKESKHIWKLQWATTILDIERSFVPVFLRKAFRSGEMVTVGKSS
DGTTPDRRWCFRVDEVNWSHWNQNLGIINEDPGKNETYQYYGFSHTVGRLRRDRWSSVVP
RVVELNKNNSNPDEVVPLDSMGNPRCDGHQQGYPRKWRTDDAPL

8/18

FIGURE 6

SEQ.ID.NO.10. Human VR3A+B+ nucleotide sequence of the coding sequence
(2229 bp).

ATGGCGGATTCCAGCGAAGGCCCGCGCGGGGGCCCGGGGAGGTGGCTGAG
CTCCCCGGGGATGAGAGTGGCACCCCAGGTGGGGAGGCTTTTCCTCTCTCCTC
CCTGGCCAATCTGTTTGAGGGGGAGGATGGCTCCCTTTCGCCCTCACCGGCTG
ATGCCAGTCGCCCTGCTGGCCCAGGCGATGGGCGACCAAATCTGCGCATGAA
GTTCCAGGGCGCCTTCCGCAAGGGGGTGCCCAACCCCATCGATCTGCTGGAG
TCCACCCTATATGAGTCCTCGGTGGTGCCTGGGCCCCAAGAAAGCACCCATGG
ACTCACTGTTTGACTACGGCACCTATCGTCACCACTCCAGTGACAACAAGAG
GTGGAGGAAGAAGATCATAGAGAAGCAGCCGCAGAGCCCCAAAGCCCCTGC
CCCTCAGCCGCCCCCCATCCTCAAAGTCTTCAACCGGCCTATCCTCTTTGACA
TCGTGTCCCGGGGCTCCACTGCTGACCTGGACGGGCTGCTCCCATTTCTTGCTG
ACCCACAAGAAACGCCTAACTGATGAGGAGTTTCGAGAGCCATCTACGGGGA
AGACCTGCCTGCCCAAGGCCTTGCTGAACCTGAGCAATGGCCGCAACGACAC
CATCCCTGTGCTGCTGGACATCGCGGAGCGCACCGGCAACATGAGGGAGTTC
ATTA ACTCGCCCTTCCGTGACATCTACTATCGAGGTCAGACAGCCCTGCACAT
CGCCATTGAGCGTCGCTGCAAACACTACGTGGA ACTTCTCGTGGCCCAGGGA
GCTGATGTCCACGCCCAGGCCCGTGGGCGCTTCTTCCAGCCCAAGGATGAGG
GGGGCTACTTCTACTTTGGGGAGCTGCCCTGTGCTGGCTGCCTGCACCAAC
CAGCCCCACATTGTCAACTACCTGACGGAGAACCCCCACAAGAAGGCGGACA
TGCGGGCGCCAGGACTCGCGAGGCAACACAGTGCTGCATGCGCTGGTGGCCAT
TGCTGACAACACCCGTGAGAACACCAAGTTTGTTACCAAGATGTACGACCTG
CTGCTGCTCAAGTGTGCCCCGCTCTTCCCCGACAGCAACCTGGAGGCCGTGCT
CAACAACGACGGCCTCTCGCCCCCTCATGATGGCTGCCAAGACGGGCAAGATT
GGGATCTTTCAGCACATCATCCGGCGGGAGGTGACGGATGAGGACACACGGC
ACCTGTCCCGCAAGTTCAAGGACTGGGCCTATGGGCCAGTGTATTCTCTCGCTT
TATGACCTCTCCTCCCTGGACACGTGTGGGGAAGAGGCCTCCGTGCTGGAGA
TCCTGGTGTACAACAGCAAGATTGAGAACCGCCACGAGATGCTGGCTGTGGA
GCCCATCAATGAACTGCTGCGGGACAAGTGGCGCAAGTTCGGGGCCGTCTCC
TTCTACATCAACGTGGTCTCCTACCTGTGTGCCATGGTCATCTTCACTCTCACC
GCCTACTACCAGCCGCTGGAGGGCACACCGCCGTACCCTTACCGCACACCGG
TGGACTACCTGCGGCTGGCTGGCGAGGTCATTACGCTCTTCACTGGGGTCTCTG
TTCTTCTTACCAACATCAAAGACTTGTTTCATGAAGAAATGCCCTGGAGTGAA
TTCTCTCTTCATTGATGGCTCCTTCCAGCTGCTCTACTTCATCTACTCTGTCCT
GGTGATCGTCTCAGCAGCCCTCTACCTGGCAGGGATCGAGGCCTACCTGGCC
GTGATGGTCTTTGCCCTGGTCTGGGCTGGATGAATGCCCTTTACTTCACCCG
TGGGCTGAAGCTGACGGGGACCTATAGCATCATGATCCAGAAGATTCTCTTC
AAGGACCTTTTCCGATTCTGCTCGTCTACTTGCTCTTCATGATCGGCTACGCT
TCAGCCCTGGTCTCCCTCCTGAACCCGTGTGCCAACATGAAGGTGTGCAATG

9/18

AGGACCAGACCAACTGCACAGTGCCCACTTACCCCTCGTGCCGTGACAGCGA
GACCTTCAGCACCTTCCTCCTGGACCTGTTTAAGCTGACCATCGGCATGGGCG
ACCTGGAGATGCTGAGCAGCACCAAGTACCCCGTGGTCTTCATCATCCTGCT
GGTGACCTACATCATCCTCACCTTTGTGCTGCTCCTCAACATGCTCATTGCCC
TCATGGGCGAGACAGTGGGCCAGGTCTCCAAGGAGAGCAAGCACATCTGGA
AGCTGCAGAGCGGCAGGCGCAGGCTGTGA

10/18
FIGURE 7

SEQ.ID.NO.11. The nucleotide sequence of human VR3A+B+ is shown including 836 bp 5' UT and 994 bp 3'UT.

TGTGCAGGCCAGGGAGGGCTTTCCAGAGGAGCCCAGTTGAGCTGGAACACCA
GTGGGGAGGAGTTGACCAGCAAAGGTGCAGGGAGGGATCAGCACTTTGCAC
TGGGGAGCAGAGTTTGTGCACTGGGGAAGTCAACTCAAGTATTGGAGCCTCA
GTTTCCTGTTCTGTAAAATGGGTTCATCATGACAGTGTGTTGATGAGGAAAAGG
ACTGCCGGCCTACACAGCAAGTCCACATGGATTTTCTGAGCCCCCTCCTGTGCC
TGAAGCCCAACGGTTAATGGTTCCTTAGCAGGTGCTTACCACGTGCCAGGC
ACTGCACTGCACTGGCCACTGGACTGCATGTTCTGTCCATGAGGCTTGGATAT
CCCCATCTTACAGATCAGGAAGCTGAGGCTATGAAATGTCGACTTGCTCAAT
GTCATGGAATGACTAAGTGTGGAGCCTGGATTTGAACTTGGCTCTCTGGGGCT
CCAAAGCTGGCTTTCTTGGTCAGCAGTAGGGTCTGGGATCCAAGTATGGGGT
CCCAGCTTGACCCTGAAGTCCACCCTCTTTCAGCTAATGCCCAAGGTAGTTGG
ACCTGGGGCCAATTTGTGTTTCCAGGTTCTGTGAAAGAGCTCCTGTTGCAGTTC
CCGCTGAGGCTTGGCGGCCAACACATCTGGGAGTGGCCTCCCTGTGCCCC
TGTCATTACAACGGTGGCTTTGAAGCAGCTGGCAGCACTGCTGCTTGTCCACG
TGGAAGGGGGCTTCCTGGAGCCCCCGCCCCTGGCCGGGTCTGCCTGACTCC
CCTTTCATTCCCTTGCAGGCTGAGCAGTGCAGACGGGCCTGGGGCAGGCATG
GCGGATTCCAGCGAAGGCCCCCCGCGCGGGGGCCCGGGGAGGTGGCTGAGCTC
CCCGGGGATGAGAGTGGCACCCCAGGTGGGGAGGCTTTTCCTCTCTCCTCCCT
GGCCAATCTGTTTGAGGGGGAGGATGGCTCCCTTTTCGCCCTCACCGGCTGATG
CCAGTCGCCCTGCTGGCCCAGGCGATGGGCGACCAAATCTGCGCATGAAGTT
CCAGGGCGCCTTCCGCAAGGGGGTGCCCAACCCCATCGATCTGCTGGAGTCC
ACCCTATATGAGTCCTCGGTGGTGCCTGGGCCCAAGAAAGCACCCATGGACT
CACTGTTTGACTACGGCACCTATCGTCACCACTCCAGTGACAACAAGAGGTG
GAGGAAGAAGATCATAGAGAAGCAGCCGCAGAGCCCCAAAGCCCCTGCCCC
TCAGCCGCCCCCATCCTCAAAGTCTTCAACCGGCCTATCCTCTTTGACATCG
TGTCCTCGGGGCTCCACTGCTGACCTGGACGGGCTGCTCCCATTTCTTGCTGACC
CACAAGAAACGCCTAACTGATGAGGAGTTTCGAGAGCCATCTACGGGGAAG
ACCTGCCTGCCCAAGGCCTTGCTGAACCTGAGCAATGGCCGCAACGACACCA
TCCCTGTGCTGCTGGACATCGCGGAGCGCACCGGCAACATGAGGGAGTTTCAT
TAACTCGCCCTTCCGTGACATCTACTATCGAGGTCAGACAGCCCTGCACATCG
CCATTGAGCGTCGCTGCAAACACTACGTGGAACTTCTCGTGGCCCAGGGAGC
TGATGTCCACGCCCAGGCCCGTGGGCGCTTCTTCCAGCCCAAGGATGAGGGG
GGCTACTTCTACTTTGGGGAGCTGCCCCTGTGCTGGCTGCCTGCACCAACCA
GCCCCACATTGTCAACTACCTGACGGAGAACCCCCACAAGAAGGCGGACATG
CGGCGCCAGGACTCGCGAGGCAACACAGTGCTGCATGCGCTGGTGGCCATTG
CTGACAACACCCGTGAGAACACCAAGTTTGTTACCAAGATGTACGACCTGCT
GCTGCTCAAGTGTGCCCCGCTCTTCCCCGACAGCAACCTGGAGGCCGTGCTC

11/18

AACAACGACGGCCTCTCGCCCCTCATGATGGCTGCCAAGACGGGCAAGATTG
GGATCTTTTCAGCACATCATCCGGCGGGAGGTGACGGATGAGGACACACGGCA
CCTGTCCCGCAAGTTCAAGGACTGGGCCTATGGGCCAGTGTATTCTCGCTTT
ATGACCTCTCCTCCCTGGACACGTGTGGGGAAGAGGCCTCCGTGCTGGAGAT
CCTGGTGTACAACAGCAAGATTGAGAACCGCCACGAGATGCTGGCTGTGGAG
CCCATCAATGAACTGCTGCGGGACAAGTGGCGCAAGTTCGGGGCCGTCTCCT
TCTACATCAACGTGGTCTCCTACCTGTGTGCCATGGTCATCTTCACTCTCACC
GCCTACTACCAGCCGCTGGAGGGGCACACCGCCGTACCCTTACCGCACACGG
TGGACTACCTGCGGCTGGCTGGCGAGGTCATTACGCTCTTCACTGGGGTCCTG
TTCTTCTTCACCAACATCAAAGACTTGTTTCATGAAGAAATGCCCTGGAGTGAA
TTCTCTCTTCATTGATGGCTCCTTCCAGCTGCTCTACTTCATCTACTCTGTCT
GGTGATCGTCTCAGCAGCCCTCTACCTGGCAGGGATCGAGGCCTACCTGGCC
GTGATGGTCTTTGCCCTGGTCCTGGGCTGGATGAATGCCCTTTACTTCACCCG
TGGGCTGAAGCTGACGGGGACCTATAGCATCATGATCCAGAAGATTCTCTTC
AAGGACCTTTTCCGATTCTGCTCGTCTACTTGCTCTTCATGATCGGCTACGCT
TCAGCCCTGGTCTCCCTCCTGAACCCGTGTGCCAACATGAAGGTGTGCAATG
AGGACCAGACCAACTGCACAGTGCCCACTTACCCCTCGTGCCGTGACAGCGA
GACCTTCAGCACCTTCCTCCTGGACCTGTTTAAGCTGACCATCGGCATGGGCG
ACCTGGAGATGCTGAGCAGCACCAAGTACCCCGTGGTCTTCATCATCCTGCT
GGTGACCTACATCATCCTCACCTTTGTGCTGCTCCTCAACATGCTCATTGCC
TCATGGGCGAGACAGTGGGCCAGGTCTCCAAGGAGAGCAAGCACATCTGGA
AGCTGCAGAGCGGCAGGCGCAGGCTGTGAGGCTCACCGATGTCCCTCCTGAC
CCTCCCTCCCCGAGTGGGCCACCACCATCCTGGACATTGAGCGCTCCTTCCC
CGTATTCTGAGGAAGGCCTTCCGCTCTGGGGAGATGGTCACCGTGGGCAAG
AGCTCGGACGGCACTCCTGACCGCAGGTGGTGCTTCAGGGTGGATGAGGTGA
ACTGGTCTCACTGGAACCAAGAACTTGGGCATCATCAACGAGGACCCGGGCAA
GAATGAGACCTACCAGTATTATGGCTTCTCGCATACCGTGGGCGCCTCCGC
AGGGATCGCTGGTCCTCGGTGGTACCCCGCGTGGTGGAAGTGAACAAGAACT
CGAACCCGGACGAGGTGGTGGTGCCTCTGGACAGCATGGGGAACCCCGCTG
CGATGGCCACCAGCAGGGTTACCCCGCAAGTGGAGGACTGATGACGCCCCG
CTCTAGGGACTGCAGCCCAGCCCCAGCTTCTCTGCCCCACTCATTTCTAGTCCA
GCCGCATTTTCAGCAGTGCCTTCTGGGGTGTCCCCCACACCCTGCTTTGGCCC
CAGAGGCGAGGGACCAGTGGAGGTGCCAGGGAGGCCCCAGGACCCTGTGGT
CCCCTGGCTCTGCCTCCCCACCCTGGGGTGGGGGCTCCCGGCCACCTGTCTTG
CTCCTATGGAGTCACATAAGCCAACGCCAGAGCCCCTCCACCTCAGGCCCCA
GCCCCTGCCTCTCCATTATTTATTTGCTCTGCTCTCAGGAAGCGACGTGACCC
CTGCCCCAGCTGGAACCTGGCAGAGGCCTTAGGACCCCGTTCCAAGTGCACT
GCCCCGCCAAGCCCCAGCCTCAGCCTGCGCCTGAGCTGCATGCGCCACCATT
TTTGGCAGCGTGGCAGCTTTGCAAGGGGCTGGGGCCCTCGGCGTGGGGCCAT
GCCTTCTGTGTGTTCTGTAGTGTCTGGGATTTGCCGGTGCTCAATAAATGTTT
ATTCATTGACGGTGGAAAAAAAAAAAAAAAAA

12/18

FIGURE 8

SEQ.ID.NO.12. Coding sequence for human VR3A+B+ (742 amino acids)

MADSSEGPRAGPGEVAELPGDESGTPGGEAFPLSSLANLFEGEDGSLSPSPADASRPAGPGDGRPNLRMKF
QGAFRKGVNPNIDLLESTLYESSVVPGPKKAPMDSLFDYGTYRHHSSDNKRWRKKIIEKQPQSPKAPAPQP
PPILKVFNRPIILFDIVSRGSTADLDGLLPFLLTHKKRLTDEEFREPSTGKTCLPKALLNLSNGRNDTIPVL
LDIAERTGNMREFINSFPRDIYYRGQTALHIAIERRCKHYVELLVAQGADVHAQARGRFFQPKDEGGYFYF
GELPLSLAACTNQPHIVNYLTENPHKKADMRRQDSRGNTVLHALVAIADNTRENTKFVTKMYDLLLLKCAR
LFPDSNLEAVLNNDGLSPLMMAAKTGKIGIFQHIIRREVTDEDTRHLSRKFKDWAYGPVYSSLYDLSSLDT
CGEEASVLEILVYNSKIENRHEMLAVEPINELLRDKWRKFGAVSFYINVVSYLCAMVIFTLTAYYQPLEGT
PPYPYRTTVDYLRRLAGEVITLFTGVLFFFTNIKDLFMKKCPGVNSLFIDGSFQLLYFIYSVLVIVSAALYL
AGIEAYLAVMVFALVLGWMNALYFTRGLKLTGTYSIMIQKILFKDLFRLLVYLLFMIGYASALVSLNPNPC
ANMKVCNEDQTNCTVPTYPSCRDSETFSTFLDLFKLTIGMGDLEMLSSTKYPVVFIIILLVTYIILTFVLL
LNMLIALMGETVGQVSKEKHIWKLQSGRRRL


13/18

FIGURE 9

	Water-injected controls	VR3 A+B-	VR3 A-B-	VR3 A+B+
Number of living oocytes	88	9	47	29
Number of dead oocytes	44	90	40	54
Percent Alive	67%	9% (* p < e - 17)	54% (p = 0.99)	35% (* p = 5.4 e-6)


14/18

FIGURE 10

a



15/18

FIGURE 10B




16/18

FIGURE 11

a VR1 and water**b****VR1 and VR3 A^+B^-** 

17/18

FIGURE 12:



18/18

FIGURE 13-

Tissue or cell type	hVR3 (mean intensity)	hVR1 (mean intensity)
Liver	900 +/- 50 (p<0.005)	55 +/- 3 (p < 0.05)
Raji lymphoma cell line	255 +/- 20 (p<0.005)	NS
Spleen	196 +/- 19 (p< 0.01)	NS
Lung	150 +/- 22 (p<0.01)	NS
DRG	129 +/- 21 (p< 0.025)	90 +/- 3 (p< 0.05)
Ovary	128 +/- 9 (p < 0.0025)	69 +/- 2 (p< 0.0005)
Placenta	120 +/- 7 (p < 0.001)	NS
Trachea	106 +/- 7 (p < 0.001)	54 +/- 4 (p< 0.01)
Small intestine	105 +/- 3 (p < 0.001)	62 +/- 5 (p< 0.01)
Prostate*	72 +/- 5 (p < 0.0025)	38 +/- 1 (p< 0.0005)
Kidney	62 +/- 4 (p < 0.05)	57 +/- 4(p< 0.005)
Spinal cord	57 +/- 2 (p< 0.00025)	47 +/- 3 (p< 0.005)

Values are the mean intensity of the labeled cRNA hybridizing to the cDNA microarray +/- S.E.M. The mean intensity for cRNAs from all tissues shown were significantly different (p value in parentheses) from 75% of the control plant cDNA value. Data are averaged from 3-6 experiments. NS: not significantly different from plant gene control (see Luo et al., 1999 for more detailed methods). * Tissue from which the VR3 was cloned.

SEQUENCE LISTING

<110> Dubin, Adrienne E
Huvar, Arne
Erlander, Mark G
Glass, Charles A

<120> DNA encoding Isoforms of the human Vanilloid Receptor
VR3

<130> Human VR3 receptors

<140>

<141>

<160> 17

<170> PatentIn Ver. 2.1

<210> 1

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 1

accggcctat cctctttgac atcgtg

26

<210> 2

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 2

2

tgtccgcctt cttgtggggg ttctc

25

<210> 3

<211> 48

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 3

aacgttggtta ccgccacat ggcggattcc agcgaaggcc cccgcgcg

48

<210> 4

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 4

taaagcggcc gcttcaggag ggacatcggt gagcctcac

39

<210> 5

<211> 2616

<212> DNA

<213> Homo sapiens

<400> 5

atggcggatt ccagcgaagg cccccgcgcg gggcccgggg aggtggctga gctccccggg 60

gatgagagtg gcacccagg tggggaggct ttctctct cctccctggc caatctgtt 120

gagggggagg atggctcct ttgccctca ccggtgatg ccagtcgcc tgcaggcca 180

ggcgatgggc gaccaaact gcgcataag ttccaggcgc cttccgcaa gggggtgcc 240

aaccctatg atctgtgga gtccacccta tatagtcct cgggtggtgc tgggccaag 300

aaagcacca tggactact gtttgactac ggcacctatc gtcaccact cagtacaac 360

aagaggtgga ggaagaagat catagagaag cagccgcaga gcccacaaagc cctgcccct 420
 cagccgcccc ccaccccaa agtcttcaac cggcctatcc tcttgacat cgtgtcccgg 480
 ggctccactg ctgacctgga cgggctgctc ccattcttgc tgaccacaa gaaacgcta 540
 actgatgagg agtttcgaga gccatctacg gggaagacct gctgcccacaa ggccttgcgtg 600
 aacctgagca atggccgcaa cgacaccatc cctgtgctgc tggacatcgc ggagcgcacc 660
 ggcaacatgc gggagttcat taactgccc ttccgtgaca tctactatcg aggtcagaca 720
 gccctgcaca tcgccattga gcgtcgtgc aaacactacg tggaaactct cgtggcccag 780
 ggagctgatg tccacgcccc gggcctggg cgcttctcc agcccaagga tgaggggggc 840

tacttctact ttggggagct gcccctgtcg ctggctgect gcaccaacca gcccacatt 900
 gtcaactacc tgacggagaa cccccacaag aaggcggaca tgcggcgcca ggactcgcga 960
 ggcaacacag tgctgcatgc gctggtggcc attgctgaca acaccgtga gaacaccaag 1020
 ttgttacca agatgtacga cctgctgctg ctcaagtgtg cccgcctctt ccccgacagc 1080
 aacctggagg ccgtgctcaa caacgacggc ctctgcccc tcatgatggc tgccaagacg 1140
 ggcaagattg ggatcttca gcacatcgc cggcgggagg tgacggatga ggacacacgg 1200
 cacctgtccc gcaagttcaa ggactgggcc tatgggccag tgtattctc gcttatgac 1260
 ctctcctccc tggacacgtg tggggaagag gcctccgtgc tggagatcct ggtgtacaac 1320
 agcaagattg agaaccgcca cgagatgctg gctgtggagc ccatcaatga actgctgcgg 1380
 gacaagtggc gcaagttcgg ggccgtctcc ttctacatca acgtggtctc ctacctgtgt 1440
 gccatggtca tcttactct caccgcctac taccagccgc tggagggcac accgccgtac 1500
 ccttaccgca ccacgggtgga ctacctgccc ctggctggcg aggtcattac gctcttact 1560
 ggggtcctgt tcttcatcac caacatcaaa gacttgttca tgaagaaatg ccctggagtg 1620
 aattctctct tcattgatgg ctcttccag ctgcttact tcatctactc tgtctggtg 1680
 atcgtctcag cagccctcta cctggcaggg atcgaggcct acctggccgt gatggtctt 1740
 gccctggtcc tgggctggat gaatgccctt tacttcccc gtgggctgaa gctgacgggg 1800
 acctatgca tcatgatcca gaagattctc ttcaaggacc tttccgatt cctgctcgtc 1860
 tacttctct tcatgatcgg ctacgttca gccctggtct cctcctgaa cccgtgtgcc 1920

aacatgaagg tgtgcaatga ggaccagacc aactgcacag tgcccactta cccctcgtgc 1980
 cgtgacagcg agacctcag cacttctc ctggacctgt ttaagctgac catcgcatg 2040
 ggcgacctgg agatgctgag cagcaccaag taccctgtgg tcttcatcat cctgctggtg 2100
 acctacatca tctcaccct tgtgctgctc ctcaacatgc tattgccct catgggcgag 2160
 acagtgggcc aggtctccaa ggagagcaag cacatctgga agctgcagtg ggccaccacc 2220
 atcctggaca ttgagcgtc ctccccgta ttctgagga aggcctccg ctctggggag 2280
 atggtcaccg tgggcaagag ctgggacggc actcctgacc gcagggtgtg cttcagggtg 2340
 gatgaggtga actggtctca ctggaaccag aacttgggca tcatcaacga ggaccgggc 2400
 aagaatgaga cctaccagta ttatggctc tcgcataccg tgggcccgcct ccgagggtat 2460
 cgtggtcct cggtggtacc ccgctgggtg gaactgaaca agaactcga cccggacgag 2520
 gtggtggtgc ctctggacag catggggaac ccccgctgcg atggccacca gcagggttac 2580
 ccccgcaagt ggaggactga tgacccccg ctctag 2616

<210> 6
<211> 3500
<212> DNA
<213> Homo sapiens

<400> 6
caattgggat ttaaaccag ggactatcca gcccacaaagc ccttcccacc acaccagggtg 60
gcctgtcctg gggccagctc tgcacacagg gcctgggtgcc cccgggggtgc ttgggaagtg 120
gcaggggcaga ggtggggccct gtggctgttc tggctcagct tctaaaacaa gagcctctgc 180
tgggggcaga gggggccgtga acccctgaaa tgttaggcag ataccctgtg ggagctttgt 240
tctgggatgc taagaaccgc ttgaggattt aagctttgcc actttggctc cggagcaagg 300
gcagaggctg agcagtgacg acgggcctgg ggcaggcatg gcggattcca gcgaaggccc 360
ccgcgcgggg cccggggagg tggctgagct cccgggggat gagagtggca cccaggtgg 420
ggaggctttt cctctctct ccttgccaa tctgtttgag ggggaggatg gctcccttc 480
gcccaccg gctgatgcca gtcgccctgc tggcccaggc gatgggcgac caaatctgcg 540
catgaagttc cagggcgctt tccgaagg ggtgcccaac cccatcgatc tctggagtc 600
caccctatat gagtctcgg tgggtgcctgg gcccagaaa gcacccatgg actcactgtt 660
tgactacggc acctatcgtc accactccag tgacaacaag aggtggagga agaagatcat 720
agagaagcag ccgcagagcc ccaaagcccc tggccctcag ccgccccca tctcaaagt 780
cttcaaccgg cctatctct ttgacatcgt gtccgggggc tccactgctg acctggacgg 840
gctgtccca ttctgtga cccacaagaa acgcctaact gatgaggagt ttgagagcc 900
atctacgggg aagacctgcc tgcccaggc cttgtgaac ctgagcaatg gccgcaacga 960
caccatccct gtgctgctgg acatcgcgga gcgcaccggc aacatgcggg agttcattaa 1020
ctgcacctc cgtgacatct actatcgagg tcagacagcc ctgcacatcg ccattgagcg 1080
tcgtgcaaaa cactacgtgg aacttctcgt ggcccaggga gctgatgtcc acggccaggc 1140
ccgtggggcg ttcttcagc ccaaggatga gggggggtac ttctactttg gggagctgcc 1200
cctgtcgtg gctgcctgca ccaaccagcc ccacattgtc aactacctga cggagaacct 1260
ccacaagaag gcggacatgc ggcgccagga ctgcgagggc aacacagtgc tgcatgcgct 1320
ggtggccatt gctgacaaca cccgtgagaa caccaagttt gttaccaaga tgtacgacct 1380
gctgtgctc aagtgtgccc gcctctccc cgacagcaac ctggaggccg tgcacaaca 1440
cgacggcctc tcggccctca tgatggctgc caagacgggc aagattggga tcttcagca 1500
catcatccgg cgggaggtga cggatgagga cacacggcac ctgtcccgca agttcaagga 1560
ctgggcctat gggccagtgt attctcgtct ttatgacctc tctccctgg acacgtgtgg 1620
ggaagaggcc tccgtgctgg agatcctggt gtacaacagc aagattgaga accgccacga 1680
gatgtggct gttggagccca tcaatgaact gctgcgggac aagtggcgca agttcggggc 1740
cgtctcctc tacatcaacg tggctccta cctgtgtgcc atggtcatct tcaactcac 1800
cgcctactac cagccgtgg agggcacacc gccgtaccct taccgacca cgggtggacta 1860
cctgcggctg gctggcgagg tcattacgt cttactggg gtctgtttct tcatcaccia 1920
catcaaagac ttgtcatga agaaatgcc tggagtgaat tctctcttca ttgatggctc 1980
cttcagctg ctctacttca tctactctgt cctggtgatc gtctcagcag cctctacct 2040
ggcagggatc gaggcctacc tggccgtgat ggtctttgcc ctggtcctgg gctggatgaa 2100

tgccctttac ttcacccgtg ggetgaagct gacggggacc tatagcatca tgatccagaa 2160
 gatttcttc aaggacctt tccgattcct gctcgtctac ttgctcttca tgatcggtca 2220
 cgcttcagcc ctggtctccc tctgaaccc gtgtgccaac atgaagggtg gcaatgagga 2280
 ccagaccaac tgcacagtgc ccacttacc ctcgtgccgt gacagcgaga ccttcagcac 2340
 ctctctctg gacctgttta agctgacct cggcatgggc gacctggaga tgetgagcag 2400
 caccaagtac cccgtggtct tcatcatcct gctggtgacc tacatcatcc tcacctttgt 2460
 gctgctctc aacatgctca ttgccctcat gggcgagaca gtggggcagg tctccaagga 2520
 gagcaagcac atctggaagc tgcagtgggc caccaccatc ctggacattg agcgctcctt 2580
 ccccgattc ctgaggaagg ccttcgctc tggggagatg gtcaccgtgg gcaagagctc 2640
 ggacggcact cctgaccgca ggtggtgctt cagggtggat gaggtgaact ggtctcactg 2700
 gaaccagaac ttgggcatca tcaacgagga ccggggcaag aatgagacct accagtatta 2760
 tggcttctcg catacgtgg gccgcctccg cagggatcgc tggctctcgg tggtaacccg 2820
 cgtggtgga aatgaacaaga actcgaaccc ggacgaggtg gtggtgcctc tggacagcat 2880
 ggggaacccc cgctcgcatg gccaccagca gggttacccc cgcaagtggg ggactgatga 2940
 cgccccgctc tagggactgc agcccagccc cagcttctct gccactcat ttctagtcca 3000
 gccgcattc agcagtgcct tctgggggtg cccccacac cctgctttgg cccagagggc 3060
 gagggaccag tggaggtgcc agggaggccc caggacctg tggccccctg gctctgcctc 3120
 cccaccttg ggtggggggt cccggccacc tgtcttctc ctatggagtc acataagcca 3180
 acgccagagc cctccacct caggccccag cccctgcctc tccattattt attgctctg 3240
 ctctcaggaa gcgacgtgac cctgccccca gctggaacct ggcagaggcc ttaggacccc 3300
 gtccaagtg cactgcccgg ccaagccccca gcctcagcct gcgcctgagc tgcctgcgcc 3360
 accatttttg gcagcgtggc agctttgcaa ggggctgggg ccctcggcgt ggggccatgc 3420
 cttctgtgtg ttctgtagt tctgggattt gccggtgctc aataaatgtt tattcattga 3480
 cgggtgaaaa aaaaaaaaaa 3500

<210> 7

<211> 871

<212> PRT

<213> Homo sapiens

<400> 7

Met Ala Asp Ser Ser Glu Gly Pro Arg Ala Gly Pro Gly Glu Val Ala

1 5 10 15

Glu Leu Pro Gly Asp Glu Ser Gly Thr Pro Gly Gly Glu Ala Phe Pro

20 25 30

Leu Ser Ser Leu Ala Asn Leu Phe Glu Gly Glu Asp Gly Ser Leu Ser

35 40 45

Pro Ser Pro Ala Asp Ala Ser Arg Pro Ala Gly Pro Gly Asp Gly Arg

50 55 60

Pro Asn Leu Arg Met Lys Phe Gln Gly Ala Phe Arg Lys Gly Val Pro
65 70 75 80

Asn Pro Ile Asp Leu Leu Glu Ser Thr Leu Tyr Glu Ser Ser Val Val
 85 90 95

Pro Gly Pro Lys Lys Ala Pro Met Asp Ser Leu Phe Asp Tyr Gly Thr
 100 105 110

Tyr Arg His His Ser Ser Asp Asn Lys Arg Trp Arg Lys Lys Ile Ile
 115 120 125

Glu Lys Gln Pro Gln Ser Pro Lys Ala Pro Ala Pro Gln Pro Pro Pro
 130 135 140

Ile Leu Lys Val Phe Asn Arg Pro Ile Leu Phe Asp Ile Val Ser Arg
145 150 155 160

Gly Ser Thr Ala Asp Leu Asp Gly Leu Leu Pro Phe Leu Leu Thr His
 165 170 175

Lys Lys Arg Leu Thr Asp Glu Glu Phe Arg Glu Pro Ser Thr Gly Lys
 180 185 190

Thr Cys Leu Pro Lys Ala Leu Leu Asn Leu Ser Asn Gly Arg Asn Asp
 195 200 205

Thr Ile Pro Val Leu Leu Asp Ile Ala Glu Arg Thr Gly Asn Met Arg
 210 215 220

Glu Phe Ile Asn Ser Pro Phe Arg Asp Ile Tyr Tyr Arg Gly Gln Thr
225 230 235 240

Ala Leu His Ile Ala Ile Glu Arg Arg Cys Lys His Tyr Val Glu Leu
 245 250 255

Leu Val Ala Gln Gly Ala Asp Val His Ala Gln Ala Arg Gly Arg Phe
 260 265 270

Phe Gln Pro Lys Asp Glu Gly Gly Tyr Phe Tyr Phe Gly Glu Leu Pro
 275 280 285

Leu Ser Leu Ala Ala Cys Thr Asn Gln Pro His Ile Val Asn Tyr Leu
290 295 300

Thr Glu Asn Pro His Lys Lys Ala Asp Met Arg Arg Gln Asp Ser Arg
305 310 315 320

Gly Asn Thr Val Leu His Ala Leu Val Ala Ile Ala Asp Asn Thr Arg
325 330 335

Glu Asn Thr Lys Phe Val Thr Lys Met Tyr Asp Leu Leu Leu Leu Lys
340 345 350

Cys Ala Arg Leu Phe Pro Asp Ser Asn Leu Glu Ala Val Leu Asn Asn
355 360 365

Asp Gly Leu Ser Pro Leu Met Met Ala Ala Lys Thr Gly Lys Ile Gly
370 375 380

Ile Phe Gln His Ile Ile Arg Arg Glu Val Thr Asp Glu Asp Thr Arg
385 390 395 400

His Leu Ser Arg Lys Phe Lys Asp Trp Ala Tyr Gly Pro Val Tyr Ser
405 410 415

Ser Leu Tyr Asp Leu Ser Ser Leu Asp Thr Cys Gly Glu Glu Ala Ser

420 425 430

Val Leu Glu Ile Leu Val Tyr Asn Ser Lys Ile Glu Asn Arg His Glu
435 440 445

Met Leu Ala Val Glu Pro Ile Asn Glu Leu Leu Arg Asp Lys Trp Arg
450 455 460

Lys Phe Gly Ala Val Ser Phe Tyr Ile Asn Val Val Ser Tyr Leu Cys
465 470 475 480

Ala Met Val Ile Phe Thr Leu Thr Ala Tyr Tyr Gln Pro Leu Glu Gly
485 490 495

Thr Pro Pro Tyr Pro Tyr Arg Thr Thr Val Asp Tyr Leu Arg Leu Ala

500 505 510
Gly Glu Val Ile Thr Leu Phe Thr Gly Val Leu Phe Phe Ile Thr Asn
515 520 525
Ile Lys Asp Leu Phe Met Lys Lys Cys Pro Gly Val Asn Ser Leu Phe
530 535 540
Ile Asp Gly Ser Phe Gln Leu Leu Tyr Phe Ile Tyr Ser Val Leu Val
545 550 555 560
Ile Val Ser Ala Ala Leu Tyr Leu Ala Gly Ile Glu Ala Tyr Leu Ala
565 570 575
Val Met Val Phe Ala Leu Val Leu Gly Trp Met Asn Ala Leu Tyr Phe
580 585 590
Thr Arg Gly Leu Lys Leu Thr Gly Thr Tyr Ser Ile Met Ile Gln Lys
595 600 605
Ile Leu Phe Lys Asp Leu Phe Arg Phe Leu Leu Val Tyr Leu Leu Phe
610 615 620
Met Ile Gly Tyr Ala Ser Ala Leu Val Ser Leu Leu Asn Pro Cys Ala
625 630 635 640
Asn Met Lys Val Cys Asn Glu Asp Gln Thr Asn Cys Thr Val Pro Thr
645 650 655
Tyr Pro Ser Cys Arg Asp Ser Glu Thr Phe Ser Thr Phe Leu Leu Asp
660 665 670
Leu Phe Lys Leu Thr Ile Gly Met Gly Asp Leu Glu Met Leu Ser Ser
675 680 685
Thr Lys Tyr Pro Val Val Phe Ile Ile Leu Leu Val Thr Tyr Ile Ile
690 695 700
Leu Thr Phe Val Leu Leu Leu Asn Met Leu Ile Ala Leu Met Gly Glu
705 710 715 720
Thr Val Gly Gln Val Ser Lys Glu Ser Lys His Ile Trp Lys Leu Gln

9

725 730 735
Trp Ala Thr Thr Ile Leu Asp Ile Glu Arg Ser Phe Pro Val Phe Leu
740 745 750
Arg Lys Ala Phe Arg Ser Gly Glu Met Val Thr Val Gly Lys Ser Ser
755 760 765
Asp Gly Thr Pro Asp Arg Arg Trp Cys Phe Arg Val Asp Glu Val Asn
770 775 780
Trp Ser His Trp Asn Gln Asn Leu Gly Ile Ile Asn Glu Asp Pro Gly
785 790 795 800
Lys Asn Glu Thr Tyr Gln Tyr Tyr Gly Phe Ser His Thr Val Gly Arg
805 810 815
Leu Arg Arg Asp Arg Trp Ser Ser Val Val Pro Arg Val Val Glu Leu
820 825 830
Asn Lys Asn Ser Asn Pro Asp Glu Val Val Val Pro Leu Asp Ser Met
835 840 845
Gly Asn Pro Arg Cys Asp Gly His Gln Gln Gly Tyr Pro Arg Lys Trp
850 855 860
Arg Thr Asp Asp Ala Pro Leu
865 870

<210> 8

<211> 2436

<212> DNA

<213> Homo sapiens

<400> 8

atggcggatt ccagcgaagg cccccgcgcg gggccccgggg aggtggctga gctccccggg 60
gatgagagtg gcaccccagg tggggagget ttctctct cctccctggc caatctgttt 120
gagggggagg atggctcct ttcgcctca cgggtgatg ccagtcgcc tgctggccca 180
ggcgtgggc gaccaaact ggcgatgaag ttccaggcg ccttcgcaa gggggtgcc 240
aaccatcg atctgctgga gtccacctat tatgagtcct cgggtgtgcc tgggccaag 300
aaagcacca tggactact gttgactac ggcacctatc gtcaccact cagtgacaac 360
aagaggtgga ggaagaagat catagagaag cagccgcaga gcccacaagc ccctgccct 420

cagccgcccc ccactctcaa agtcttcaac cggcctatcc tctttgacat cgtgtcccgg 480
 ggctccactg ctgacctgga cgggctgctc ccattcttgc tgaccacaaa gaaacgcta 540
 actgatgagg agtttcgaga gccatctacg gggaagacct gcctgccccaa ggccttgctg 600
 aacctgagca atggccgcaa cgacaccatc cctgtgctgc tggacatcgc ggagcgcacc 660
 ggcaacatgc gggagttcat taactgcccc ttccgtgaca tctactatcg aggtcagaca 720
 gccctgcaca tcgccattga gcgtcgtctc aaacactacg tggaaactct cgtggcccag 780
 ggagctgatg tccacgcccc ggcccgtggg cgcttcttcc agcccaagga tgaggggggc 840
 tactctact ttggggagct gcccctgtcg ctggctgcct gcaccaacca gcccacatt 900
 gtcaactacc tgacggagaa cccccacaag aaggcggaca tgcggcgcca ggactcgcga 960
 ggcaacacag tgctgcacgc gctgggtggc attgctgaca acaccgtga gaacaccaag 1020
 ttgttacca agatgtacga cctgtgctg ctcaagtgtg cccgcctct ccccgacagc 1080

aacctggagg cctgtctcaa caacgacggc ctctgcccc tcatgatggc tgccaagacg 1140
 ggcaagattg agaaccgcca cgagatgctg gctgtggagc ccataatga actgctgcgg 1200
 gacaagtggc gcaagttcgg ggccgtctcc ttctacatca acgtggtctc ctacctgtgt 1260
 gccatggtea tcttactct caccgcctac taccagccgc tggagggcac accgccgtac 1320
 ccttaccgca ccacggtgga ctacctgcgg ctggctggcg aggtcattac gctcttact 1380
 ggggtcctgt tcttcatcac caacatcaaa gacttgttca tgaagaaatg cctggagtgt 1440
 aattctctct tcattgatgg ctcttccag ctgctctact tcactactc tctctgggtg 1500
 atcgtctcag cagccctcta cctggcaggg atcgaggcct acctggccgt gatggtcttt 1560
 gccctggctc tgggtgggat gaatgccctt tacttcaccc gtgggctgaa gctgacgggg 1620
 acctatagca tcatgatcca gaagattctc ttcaaggacc ttcccgatt cctgctcgtc 1680
 tacttctct tcatgatcgg ctacgttca gccctgggtct cctctctgaa cccgtgtgcc 1740
 aacatgaagg tgtgcaatga ggaccagacc aactgcacag tgcccactta cccctcgtgc 1800
 cgtgacagcg agacctcag caccttctc ctggacctgt ttaagctgac catcggcacg 1860
 ggagacctgg agatgctgag cagcaccaag taccctgtgg tcttcatcat cctgctggtg 1920
 acctacatca tctcaccct tgtgtgctc ctcaacatgc tcattgccct catgggcgag 1980
 acagtggggc aggtctccaa ggagagcaag cacatctgga agctgcagtg ggccaccacc 2040
 atcctggaca ttgagcgctc ctccccgta ttctgagga aggccttccg ctctggggag 2100
 atggtcaccg tgggcaagag ctccgacggc actcctgacc gcaggtgggtg cttcaggggtg 2160
 gatgaggtga actggttca ctggaaccag aacttgggca tcatcaacga ggaccgggc 2220
 agaatgaga cctaccagta ttatggcttc tcgcataacc tgggccgcct ccgcagggat 2280
 cgctggtcct cgttggtacc ccgcgtggtg gaactgaaca agaactcgaa cccggacgag 2340
 gtggtggtgc ctctggacag catggggaac ccccgctgcg atggccacca gcagggttac 2400
 ccccgcaagt ggaggactga tgacccccg ctctag 2436

<210> 9

<211> 811

<212> PRT

<213> Homo sapiens

11

<400> 9

Met Ala Asp Ser Ser Glu Gly Pro Arg Ala Gly Pro Gly Glu Val Ala
 1 5 10 15

Glu Leu Pro Gly Asp Glu Ser Gly Thr Pro Gly Gly Glu Ala Phe Pro
 20 25 30

Leu Ser Ser Leu Ala Asn Leu Phe Glu Gly Glu Asp Gly Ser Leu Ser
 35 40 45

Pro Ser Pro Ala Asp Ala Ser Arg Pro Ala Gly Pro Gly Asp Gly Arg
 50 55 60

Pro Asn Leu Arg Met Lys Phe Gln Gly Ala Phe Arg Lys Gly Val Pro
 65 70 75 80

Asn Pro Ile Asp Leu Leu Glu Ser Thr Leu Tyr Glu Ser Ser Val Val
 85 90 95

Pro Gly Pro Lys Lys Ala Pro Met Asp Ser Leu Phe Asp Tyr Gly Thr
 100 105 110

Tyr Arg His His Ser Ser Asp Asn Lys Arg Trp Arg Lys Lys Ile Ile
 115 120 125

Glu Lys Gln Pro Gln Ser Pro Lys Ala Pro Ala Pro Gln Pro Pro Pro
 130 135 140

Ile Leu Lys Val Phe Asn Arg Pro Ile Leu Phe Asp Ile Val Ser Arg
 145 150 155 160

Gly Ser Thr Ala Asp Leu Asp Gly Leu Leu Pro Phe Leu Leu Thr His
 165 170 175

Lys Lys Arg Leu Thr Asp Glu Glu Phe Arg Glu Pro Ser Thr Gly Lys
 180 185 190

Thr Cys Leu Pro Lys Ala Leu Leu Asn Leu Ser Asn Gly Arg Asn Asp
 195 200 205

Thr Ile Pro Val Leu Leu Asp Ile Ala Glu Arg Thr Gly Asn Met Arg

12

210 215 220
Glu Phe Ile Asn Ser Pro Phe Arg Asp Ile Tyr Tyr Arg Gly Gln Thr
225 230 235 240
Ala Leu His Ile Ala Ile Glu Arg Arg Cys Lys His Tyr Val Glu Leu
245 250 255
Leu Val Ala Gln Gly Ala Asp Val His Ala Gln Ala Arg Gly Arg Phe
260 265 270
Phe Gln Pro Lys Asp Glu Gly Gly Tyr Phe Tyr Phe Gly Glu Leu Pro
275 280 285
Leu Ser Leu Ala Ala Cys Thr Asn Gln Pro His Ile Val Asn Tyr Leu
290 295 300
Thr Glu Asn Pro His Lys Lys Ala Asp Met Arg Arg Gln Asp Ser Arg
305 310 315 320
Gly Asn Thr Val Leu His Ala Leu Val Ala Ile Ala Asp Asn Thr Arg
325 330 335
Glu Asn Thr Lys Phe Val Thr Lys Met Tyr Asp Leu Leu Leu Leu Lys
340 345 350
Cys Ala Arg Leu Phe Pro Asp Ser Asn Leu Glu Ala Val Leu Asn Asn
355 360 365
Asp Gly Leu Ser Pro Leu Met Met Ala Ala Lys Thr Gly Lys Ile Glu
370 375 380
Asn Arg His Glu Met Leu Ala Val Glu Pro Ile Asn Glu Leu Leu Arg
385 390 395 400
Asp Lys Trp Arg Lys Phe Gly Ala Val Ser Phe Tyr Ile Asn Val Val
405 410 415
Ser Tyr Leu Cys Ala Met Val Ile Phe Thr Leu Thr Ala Tyr Tyr Gln
420 425 430

Pro Leu Glu Gly Thr Pro Pro Tyr Pro Tyr Arg Thr Thr Val Asp Tyr
435 440 445

Leu Arg Leu Ala Gly Glu Val Ile Thr Leu Phe Thr Gly Val Leu Phe
450 455 460

Phe Ile Thr Asn Ile Lys Asp Leu Phe Met Lys Lys Cys Pro Gly Val
465 470 475 480

Asn Ser Leu Phe Ile Asp Gly Ser Phe Gln Leu Leu Tyr Phe Ile Tyr
485 490 495

Ser Val Leu Val Ile Val Ser Ala Ala Leu Tyr Leu Ala Gly Ile Glu
500 505 510

Ala Tyr Leu Ala Val Met Val Phe Ala Leu Val Leu Gly Trp Met Asn
515 520 525

Ala Leu Tyr Phe Thr Arg Gly Leu Lys Leu Thr Gly Thr Tyr Ser Ile
530 535 540

Met Ile Gln Lys Ile Leu Phe Lys Asp Leu Phe Arg Phe Leu Leu Val
545 550 555 560

Tyr Leu Leu Phe Met Ile Gly Tyr Ala Ser Ala Leu Val Ser Leu Leu
565 570 575

Asn Pro Cys Ala Asn Met Lys Val Cys Asn Glu Asp Gln Thr Asn Cys
580 585 590

Thr Val Pro Thr Tyr Pro Ser Cys Arg Asp Ser Glu Thr Phe Ser Thr
595 600 605

Phe Leu Leu Asp Leu Phe Lys Leu Thr Ile Gly Met Gly Asp Leu Glu
610 615 620

Met Leu Ser Ser Thr Lys Tyr Pro Val Val Phe Ile Ile Leu Leu Val
625 630 635 640

Thr Tyr Ile Ile Leu Thr Phe Val Leu Leu Leu Asn Met Leu Ile Ala
645 650 655

Leu Met Gly Glu Thr Val Gly Gln Val Ser Lys Glu Ser Lys His Ile

14

660 665 670

Trp Lys Leu Gln Trp Ala Thr Thr Ile Leu Asp Ile Glu Arg Ser Phe
 675 680 685

Pro Val Phe Leu Arg Lys Ala Phe Arg Ser Gly Glu Met Val Thr Val
 690 695 700

Gly Lys Ser Ser Asp Gly Thr Pro Asp Arg Arg Trp Cys Phe Arg Val
 705 710 715 720

Asp Glu Val Asn Trp Ser His Trp Asn Gln Asn Leu Gly Ile Ile Asn
 725 730 735

Glu Asp Pro Gly Lys Asn Glu Thr Tyr Gln Tyr Tyr Gly Phe Ser His
 740 745 750

Thr Val Gly Arg Leu Arg Arg Asp Arg Trp Ser Ser Val Val Pro Arg
 755 760 765

Val Val Glu Leu Asn Lys Asn Ser Asn Pro Asp Glu Val Val Val Pro
 770 775 780

Leu Asp Ser Met Gly Asn Pro Arg Cys Asp Gly His Gln Gln Gly Tyr
 785 790 795 800

Pro Arg Lys Trp Arg Thr Asp Asp Ala Pro Leu
 805 810

<210> 10

<211> 2229

<212> DNA

<213> Homo sapiens

<400> 10

atggcggatt ccagcgaagg cccccgcgcg gggcccgagg aggtggctga gctccccggg 60
 gatgagagtg gcaccccagg tggggaggct ttctctct cctccctggc caatctgtt 120
 gagggggagg atggctccct ttgccctca ccggctgatg ccagtcgcc tgctggccca 180
 ggcgatgggc gaccaaactc gcgcatgaag ttccaggcgc cttccgcaa gggggtgccc 240
 aaccccatcg atctgctgga gtccacccta tatgagtct cgggtgtgcc tgggccaag 300
 aaagcaccca tggactact gttgactac ggcacctatc gtcaccactc cagtgacaac 360

aagaggtgga ggaagaagat catagagaag cagccgcaga gccccaaagc ccctgccct 420
cagccgcccc ccacctcaa agtcttcaac cggcctatcc tctttgacat cgtgtcccgg 480
ggctccactg ctgacctgga cgggctgctc ccattcttgc tgaccacaa gaaacgccta 540
actgatgagg agtttcgaga gccatctacg gggaagacct gcctgccccaa ggccttgctg 600
aacctgagca atggccgcaa cgacaccatc cctgtgctgc tggacatgc ggagcgcacc 660
ggcaacatga gggagtcat taactgccc ttcctgaca tctactatcg aggtcagaca 720
gccctgcaca tcgccattga gcgtcgtgc aaacactacg tggaaactct cgtggccag 780
ggagctgatg tccacgcccc ggcccgtggg cgcttcttc agcccaagga tgaggggggc 840
tacttctact ttggggagct gccctgtcg ctggctgcct gcaccaacca gcccacatt 900
gtcaactacc tgacggagaa cccccacaag aaggcggaca tgcggcgcca ggactcgcga 960
ggcaacacag tctgcatgc gctggtggcc attgtgaca acaccgtga gaacaccaag 1020
ttgttacca agatgtacga cctgctgctg ctcaagtgtg cccgcctctt ccccgacagc 1080
aacctggagg ccgtgctcaa caacgacggc ctctgcccc tcatgatggc tgccaagacg 1140
ggcaagattg ggatcttca gcacatcatc cggcgggagg tgacggatga ggacacacgg 1200
cacctgtccc gcaagtcaa ggactgggccc tatgggccag tgtattctc gctttatgac 1260
ctctctccc tggacacgtg tggggaagag gcctccgtgc tggagatcct ggtgtacaac 1320
agcaagattg agaaccgcca cgagatgctg gctgtggagc ccatcaatga actgctgcgg 1380
gacaagtggc gcaagtccgg ggccgtctcc ttctacatca acgtggtctc ctacctgtg 1440
gccatggtca tcttactct caccgcctac taccagccgc tggagggcac accgcctac 1500
ccttaccgca ccacgggtgga ctacctgcgg ctggctggcg aggtcattac gctcttact 1560
ggggtcctgt tcttctac caacatcaaa gactgttca tgaagaaatg ccctggagtg 1620
aattctctct tcattgatgg ctcttccag ctgcttact tcatctactc tgcctggtg 1680
atcgtctcag cagccctcta cctggcaggg atcgaggcct acctggcctg gatggtctt 1740
gccctggctc tgggctggat gaatgccctt tacttcacc gtgggctgaa gctgacgggg 1800

acctatagca tcatgatcca gaagattctc ttcaaggacc ttctcgatt cctgctcgtc 1860
tacttctct tcatgatcgg ctacgttca gccctggtct cctctctgaa cccgtgtgcc 1920
aacatgaagg tgtgcaatga ggaccagacc aactgcacag tgcccactta ccctcgtgc 1980
cgtgacagcg agacctcag caccttctc ctggacctgt ttaagctgac catcggcag 2040
ggcgacctgg agatgctgag cagcaccaag taccctgtgg tcttcatcat cctgctggtg 2100
acctacatca tcttacctt tgtctgctc ctcaacatgc tcattgccct catgggcgag 2160
acagtgggcc aggtctccaa ggagagcaag cacatctgga agctgcagag cggcaggcgc 2220
aggctgtga 2229

<210> 11

<211> 4059

<212> DNA

<213> Homo sapiens

<400> 11

tgtgcaggcc agggagggct ttccagagga gcccagttga gctggaacac cagtggggag 60

gagttgacca gcaaagggtgc agggaggggat cagcactttg cactggggag cagagtttgt 120
gcactgggga agtcaactca agtattggag cctcagtttc ctgttctgta aaatgggttc 180
atcatgacag tgtttgatga ggaaaaggac tgccggccta cacagcaagt ccacatggat 240
tttctgagcc cctcctgtgc ctgaagccca cggttaatgg ttctgcctta gcaggtgctt 300
accacgtgcc aggcactgca ctgcactggc cactggactg catgttctgt ccatgaggt 360
tggatatccc catcttacag atcaggaagc tgaggctatg aaatgtcgac ttgctcaatg 420
tcattggaatg actaagtgtg gagcctggat ttgaacttgg ctctctgggg ctccaaagct 480
ggctttcttg gtcagcagta gggctctggga tccaagtatg ggggtccagc ttgacctga 540
agtccacctt ctttcagcta atgccaagg tagttggacc tggggccaat ttgttttcc 600
agggtcgtga aagagctcct gttgcagttc ccgcctgagg cttggcggcc aaccacatct 660
gggagtgccc tcctgtgcc cctgtcatta caacgggtggc ttgaagcag ctggcagcac 720
tgctgcttgt ccacgtggaa gggggcttcc tggagcccc gccctggcc gggttctgcc 780
tgactcccc ttattccct tcaggtctga gcagtgcaga cgggcctggg gcaggcatgg 840
cggattccag cgaaggcccc cgcgcggggc cgggggagggt ggctgagctc cccggggatg 900
agagtggcac ccaggtggg gaggttttc ctctctctc cctggccaat ctgtttgagg 960
gggaggatgg ctcccttctg cctcaccgg ctgatgccag tcgccctgct ggcccaggcg 1020
atgggcgacc aaatctgcgc atgaagtcc agggcgctt ccgaagggg gtgccaacc 1080
ccatcgatct gctggagtc accctatatg agtcctcgtt ggtgcctggg cccaagaaag 1140
cacccatgga ctactgtt gactacggca cctatcgta ccactccagt gacaacaaga 1200
gggtggaggaa gaagatcata gagaagcagc cgcagagccc caaagccct gccctcagc 1260
cgcccccat cctcaaagtc ttcaaccggc ctatctctt tgacatcgtg tcccggggct 1320
ccactgctga cctggacggg ctgtcccat tcttctgac ccacaagaaa cgcctaactg 1380
atgaggagtt tcgagagcca tctacgggga agacctgctt gcccaaggcc ttgctgaacc 1440
tgagcaatgg ccgaacgac accatccctg tgctgctgga catcgcgag cgcaccggca 1500
acatgaggga gttcattaac tcgcccttc gtgacatcta ctatcgaggt cagacagccc 1560
tgcacatgc cattgagcgt cgctgcaaac actacgtgga acttctctg gcccaggag 1620
ctgatgtcca cgcccaggcc cgtgggcgt tcttcagcc caaggatgag gggggctact 1680
tctactttgg ggagctgccc ctgtcgtgg ctgctgcac caaccagccc cacattgtca 1740
actacctgac ggagaacccc cacaagaagg cggacatgcg gcgccaggac tcgcgaggca 1800
acacagtgt gcatgcgctg gtggccattg ctgacaacac ccgtgagaac accaagtttg 1860
ttaccaagat gtacgacctg ctgctgctca agtgtgccc cctctcccc gacagcaacc 1920
tggaggccgt gctcaacaac gacggcctct cgcctctcat gatggctgcc aagacgggca 1980
agattgggat ctttcagcac atcatcggc gggaggtgac ggatgaggac acacggcacc 2040
tgtcccgcaa gttcaaggac tgggcctatg ggccagtgtg ttctcgtt tatgacctt 2100
cctccctgga cagtggtggg gaagaggcct ccgtgctgga gatctggtg tacaacagca 2160
agattgagaa ccgccacgag atgctggctg tggagcccat caatgaactg ctgcgggaca 2220
agtggcgcaa gttcggggcc gtctcttct acatcaacgt ggtctctac ctgtgtgcca 2280
tggctatct cactctacc gcctactacc agccgtgga gggcacacc ccgtaccct 2340

accgcaccac ggtggactac ctgcggctgg ctggcgaggt cattacgtc ttactgggg 2400
tctgttctt cttaccaac atcaaagact tgtcatgaa gaaatgccct ggagtgaatt 2460
ctcttctcat tgatggctcc ttccagctgc tctacttcat ctactctgc ctggtgatc 2520

tctcagcagc cctctacctg gcagggatcg aggcctacct ggccgtgatg gtctttgccc 2580
 tggctcctggg ctggatgaat gccctttact tcacccgtgg gctgaagctg acggggacct 2640
 atagcatcat gatccagaag attctcttca aggacctttt ccgattcctg ctcgtctact 2700
 tgctcttcat gatcggctac gcttcagccc tggctcctct cctgaaccgg tgtgccaaca 2760
 tgaaggtgtg caatgaggac cagaccaact gcacagtgcc cacttaccct tctgcccgtg 2820
 acagcgagac cttcagcacc ttctcctgg accgttttaa gctgaccatc ggcatgggcg 2880
 acctggagat gctgagcagc accaagtacc ccgtgggtct catcatcctg ctggtgacct 2940
 acatcatcct cacctttgtg ctgctctca acatgctcat tgcctcatg ggcgagacag 3000
 tgggccaggt ctccaaggag agcaagcaca tctggaagct gcagagcggc aggcgcaggc 3060
 tgtgaggctc accgatgtcc ctctgaccc tccctccccg cagtgggcca ccaccatct 3120
 ggacattgag cgtctctcc ccgtattct gaggaaggcc ttccgtctg gggagatggt 3180
 caccgtgggc aagagctcgg acggcactcc tgaccgcagg tgggtgctca ggtggtatga 3240
 ggtgaactgg tctcactgga accagaactt gggcatcatc aacgaggacc cgggcaagaa 3300
 tgagacctac cagtattatg gcttctcgca taccgtgggc cgcctccgca gggatcgctg 3360
 gtctcctggt gtaccccgcg tgggtggaact gaacaagaac tgaaccggc acgaggtggt 3420
 ggtgcctctg gacagcatgg ggaacccccg ctgcgatggc caccagcagg gttacccccg 3480
 caagtggagg actgatgacg ccccgctcta gggactgcag cccagcccca gtttctctgc 3540
 ccactcattt ctagtccagc cgcatttcag cagtgccttc tggggtgtcc cccacaccc 3600
 tgctttggcc ccagaggcga gggaccagtg gaggtgccag ggaggcccca ggacctgtg 3660
 gtccccggc tctgcctccc caccctgggg tggggggtcc cggccactg tcttctct 3720
 atggagtcac ataagccaac gccagagccc ctccacctca ggccccagcc cctgcctctc 3780
 cattatttat ttgtctgtct ctcaggaagc gacgtgaccc ctgccccagc tggaacctgg 3840
 cagaggcctt aggaccccg tccaagtgc ctgcccggcc aagccccagc ctcagcctgc 3900
 gcctgagctg catgcgccac ctttttggc agcgtggcag ctttgcaagg ggctggggcc 3960
 ctggcgctgg ggccatgcct tctgtgtgt ctgtagtgtc tgggatttgc cggtgctcaa 4020
 taaatgttta ttcattgacg gtggaaaaaa aaaaaaaaaa 4059

<210> 12

<211> 742

<212> PRT

<213> Homo sapiens

<400> 12

Met Ala Asp Ser Ser Glu Gly Pro Arg Ala Gly Pro Gly Glu Val Ala
 1 5 10 15

Glu Leu Pro Gly Asp Glu Ser Gly Thr Pro Gly Gly Glu Ala Phe Pro
 20 25 30

Leu Ser Ser Leu Ala Asn Leu Phe Glu Gly Glu Asp Gly Ser Leu Ser
 35 40 45

18

Pro Ser Pro Ala Asp Ala Ser Arg Pro Ala Gly Pro Gly Asp Gly Arg
50 55 60

Pro Asn Leu Arg Met Lys Phe Gln Gly Ala Phe Arg Lys Gly Val Pro
65 70 75 80

Asn Pro Ile Asp Leu Leu Glu Ser Thr Leu Tyr Glu Ser Ser Val Val
85 90 95

Pro Gly Pro Lys Lys Ala Pro Met Asp Ser Leu Phe Asp Tyr Gly Thr
100 105 110

Tyr Arg His His Ser Ser Asp Asn Lys Arg Trp Arg Lys Lys Ile Ile
115 120 125

Glu Lys Gln Pro Gln Ser Pro Lys Ala Pro Ala Pro Gln Pro Pro Pro
130 135 140

Ile Leu Lys Val Phe Asn Arg Pro Ile Leu Phe Asp Ile Val Ser Arg
145 150 155 160

Gly Ser Thr Ala Asp Leu Asp Gly Leu Leu Pro Phe Leu Leu Thr His
165 170 175

Lys Lys Arg Leu Thr Asp Glu Glu Phe Arg Glu Pro Ser Thr Gly Lys
180 185 190

Thr Cys Leu Pro Lys Ala Leu Leu Asn Leu Ser Asn Gly Arg Asn Asp
195 200 205

Thr Ile Pro Val Leu Leu Asp Ile Ala Glu Arg Thr Gly Asn Met Arg
210 215 220

Glu Phe Ile Asn Ser Pro Phe Arg Asp Ile Tyr Tyr Arg Gly Gln Thr
225 230 235 240

Ala Leu His Ile Ala Ile Glu Arg Arg Cys Lys His Tyr Val Glu Leu
245 250 255

Leu Val Ala Gln Gly Ala Asp Val His Ala Gln Ala Arg Gly Arg Phe
260 265 270

Phe Gln Pro Lys Asp Glu Gly Gly Tyr Phe Tyr Phe Gly Glu Leu Pro
275 280 285

Leu Ser Leu Ala Ala Cys Thr Asn Gln Pro His Ile Val Asn Tyr Leu
290 295 300

Thr Glu Asn Pro His Lys Lys Ala Asp Met Arg Arg Gln Asp Ser Arg
305 310 315 320

Gly Asn Thr Val Leu His Ala Leu Val Ala Ile Ala Asp Asn Thr Arg
325 330 335

Glu Asn Thr Lys Phe Val Thr Lys Met Tyr Asp Leu Leu Leu Leu Lys
340 345 350

Cys Ala Arg Leu Phe Pro Asp Ser Asn Leu Glu Ala Val Leu Asn Asn
355 360 365

Asp Gly Leu Ser Pro Leu Met Met Ala Ala Lys Thr Gly Lys Ile Gly
370 375 380

Ile Phe Gln His Ile Ile Arg Arg Glu Val Thr Asp Glu Asp Thr Arg
385 390 395 400

His Leu Ser Arg Lys Phe Lys Asp Trp Ala Tyr Gly Pro Val Tyr Ser
405 410 415

Ser Leu Tyr Asp Leu Ser Ser Leu Asp Thr Cys Gly Glu Glu Ala Ser
420 425 430

Val Leu Glu Ile Leu Val Tyr Asn Ser Lys Ile Glu Asn Arg His Glu
435 440 445

Met Leu Ala Val Glu Pro Ile Asn Glu Leu Leu Arg Asp Lys Trp Arg
450 455 460

Lys Phe Gly Ala Val Ser Phe Tyr Ile Asn Val Val Ser Tyr Leu Cys
465 470 475 480

20

Ala Met Val Ile Phe Thr Leu Thr Ala Tyr Tyr Gln Pro Leu Glu Gly
485 490 495

Thr Pro Pro Tyr Pro Tyr Arg Thr Thr Val Asp Tyr Leu Arg Leu Ala
500 505 510

Gly Glu Val Ile Thr Leu Phe Thr Gly Val Leu Phe Phe Phe Thr Asn
515 520 525

Ile Lys Asp Leu Phe Met Lys Lys Cys Pro Gly Val Asn Ser Leu Phe
530 535 540

Ile Asp Gly Ser Phe Gln Leu Leu Tyr Phe Ile Tyr Ser Val Leu Val
545 550 555 560

Ile Val Ser Ala Ala Leu Tyr Leu Ala Gly Ile Glu Ala Tyr Leu Ala
565 570 575

Val Met Val Phe Ala Leu Val Leu Gly Trp Met Asn Ala Leu Tyr Phe
580 585 590

Thr Arg Gly Leu Lys Leu Thr Gly Thr Tyr Ser Ile Met Ile Gln Lys
595 600 605

Ile Leu Phe Lys Asp Leu Phe Arg Phe Leu Leu Val Tyr Leu Leu Phe
610 615 620

Met Ile Gly Tyr Ala Ser Ala Leu Val Ser Leu Leu Asn Pro Cys Ala
625 630 635 640

Asn Met Lys Val Cys Asn Glu Asp Gln Thr Asn Cys Thr Val Pro Thr
645 650 655

Tyr Pro Ser Cys Arg Asp Ser Glu Thr Phe Ser Thr Phe Leu Leu Asp
660 665 670

Leu Phe Lys Leu Thr Ile Gly Met Gly Asp Leu Glu Met Leu Ser Ser
675 680 685

Thr Lys Tyr Pro Val Val Phe Ile Ile Leu Leu Val Thr Tyr Ile Ile

21

690 695 700

Leu Thr Phe Val Leu Leu Leu Asn Met Leu Ile Ala Leu Met Gly Glu
705 710 715 720

Thr Val Gly Gln Val Ser Lys Glu Ser Lys His Ile Trp Lys Leu Gln
725 730 735

Ser Gly Arg Arg Arg Leu
740

<210> 13
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 13
ctacctgacg gagaaccccc acaag 25

<210> 14
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 14
gtagtaggcg gtgagactga agatga 26

<210> 15
<211> 51
<212> DNA
<213> Artificial Sequence

22

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 15

aacgttggcg gccgcgccac catggcggat tccagcgaag gccccgcgc g 51

<210> 16

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 16

aacgtttcta gactgggctg cagtcctag 30

<210> 17

<211> 233

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: DNA MicroArray
probe

<400> 17

ccaccatcct ggacattgag cgctccttcc ccgtattcct gaggaaggcc ttccgctctg 60
gggagatggt caccgtgggc aagagctcgg acggcactcc tgaccgcagt ggtgcttcag 120
ggtggatgag gtgaactggt ctactggaa ccagaacttg ggcatcatca acgaggacc 180
gggcaagaat gagacctacc agtattatgg cttctcgcat accgtgggcc gcc 233

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/03456

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1, 23.5; 530/350, 387.1, 387.9, 388.1; 435/7.1, 7.2, 69.1, 71.1, 71.2, 325, 252.3, 254.11, 320.1; 514/2, 8, 12

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99/37765 A1 (SMITHKLINE BEECHAM PLC) 29 July 1999 (29/07/99), see entire document.	1-22



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

21 MAY 2001

Date of mailing of the international search report

08 JUN 2001

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PREMA MERTZ

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/03456

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

C07K 14/47, 14/705, 14/72, 16/18, 16/22, 16/26; G01N 33/53, 33/567; C12N 5/10, 15/12, 15/63, 15/64

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

536/23.1, 23.5; 530/350, 387.1, 387.9, 388.1; 435/7.1, 7.2, 69.1, 71.1, 71.2, 325, 252.3, 254.11, 320.1; 514/2, 8, 12

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

WEST, CAS ONLINE, MEDLINE, CAPLUS

search terms: human VR3 receptor, nucleic acid, polynucleotide, DNA, recombinant, antibody, assay, agonists, antagonists, method, tehrapy, treatment, administer